

Bonzo Antibody
Catalog # ASC10033

Specification

Bonzo Antibody - Product Information

Application	WB
Primary Accession	O00574
Other Accession	AAB64221 , 2253422
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	Bonzo antibody can be used for detection of Bonzo by Western blot at 1:1000 dilution.

Bonzo Antibody - Additional Information

Gene ID **10663**

Other Names

Bonzo Antibody: BONZO, CD186, STRL33, TYMSTR, BONZO, C-X-C chemokine receptor type 6, CDw186, CXC-R6, chemokine (C-X-C motif) receptor 6

Target/Specificity

CXCR6;

Reconstitution & Storage

Bonzo antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

Bonzo Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Bonzo Antibody - Protein Information

Name CXCR6

Synonyms BONZO, STRL33, TYMSTR

Function

Receptor for the C-X-C chemokine CXCL16. Used as a coreceptor by SIVs and by strains of HIV-2 and m-tropic HIV-1.

Cellular Location

Cell membrane; Multi-pass membrane protein.

Tissue Location


Expressed in lymphoid tissues and activated T cells

Bonzo Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Bonzo Antibody - Images



Western blot analysis of Bonzo in SW1353 total cells lysate with Bonzo antibody at 1:1000 dilution.

Bonzo Antibody - Background

Bonzo Antibody: Human immunodeficiency virus (HIV) and simian immunodeficiency virus (SIV) require coreceptors, in addition to CD4, to infect target cells. Some G protein-coupled receptors including CCR5, CXCR4, CCR3, and CCR2b in the chemokine receptor family have been identified as HIV coreceptors. An orphan G protein-coupled receptor was recently cloned and designated Bonzo, STRL33 and TYMSTR, and identified as HIV and SIV coreceptor. Bonzo/STRL33 is used by SIV, HIV-2 and HIV-1. The messenger RNA of Bonzo/STRL33 is expressed in lymphoid tissues and activated peripheral blood lymphocytes.

Bonzo Antibody - References

- Deng HK, Unutmaz D, KewalRamani VN, Littman DR. Expression cloning of new receptors used by simian and human immunodeficiency viruses. *Nature* 1997;388:296-300
- Liao F, Alkhatib G, Peden KW, Sharma G, Berger EA, Farber JM. STRL33, A novel chemokine receptor-like protein, functions as a fusion cofactor for both macrophage-tropic and T cell line-tropic HIV-1. *J Exp Med* 1997;185:2015-23
- Alkhatib G, Liao F, Berger EA, Farber JM, Peden KW. A new SIV co-receptor, STRL33. *Nature* 1997;388:238
- Loetscher M, Amara A, Oberlin E, Brass N, Legler D, Loetscher P, D'Apuzzo M, Meese E, Rousset D, Virelizier JL, Baggiolini M, Arenzana-Seisdedos F, Moser B. TYMSTR, a putative chemokine receptor selectively expressed in activated T cells, exhibits HIV-1 coreceptor function. *Curr Biol* 1997;7:652-60 (RD1299)