

Noxa Antibody
Catalog # ASC10137**Specification**

Noxa Antibody - Product Information

Application	WB, IHC, IF
Primary Accession	O9JM54
Other Accession	NP_067426 , 10946832
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	Noxa antibody can be used for detection of Noxa by Western blot at at 0.5 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 1 µg/mL. For immunofluorescence start at 10 µg/mL.

Noxa Antibody - Additional Information

Gene ID 58801

Other Names

Noxa Antibody: Noxa, Phorbol-12-myristate-13-acetate-induced protein 1, phorbol-12-myristate-13-acetate-induced protein 1

Target/Specificity

Pmaip1;

Reconstitution & Storage

Noxa antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

Noxa Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Noxa Antibody - Protein Information

Name Pmaip1

Synonyms Noxa

Function

Promotes activation of caspases and apoptosis. Promotes mitochondrial membrane changes and efflux of apoptogenic proteins from the mitochondria. Contributes to p53/TP53-dependent apoptosis after radiation exposure. Promotes proteasomal degradation of MCL1. Competes with BIM/BCL2L11 for binding to MCL1 and can displace BIM/BCL2L11 from its binding site on MCL1 (By similarity). Competes with BAK1 for binding to MCL1 and can displace BAK1 from its binding site

on MCL1.

Cellular Location

Mitochondrion.

Tissue Location


Detected in thymocytes after irradiation with X- rays. Not detectable in untreated thymocytes (at protein level) Detected in embryonic neural precursor cells of the telencephalon Constitutively expressed at low levels in adult brain, testis, thymus, spleen, lung and kidney.

Noxa Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Noxa Antibody - Images



Western blot analysis of Noxa in human stomach tissue lysate with Noxa antibody at (A) 0.5, (B) 1 and (C) 2 µg/mL.



Immunohistochemistry of Noxa in human stomach tissue with Noxa antibody at 1 µg/mL.



Immunofluorescence of Noxa in Human Stomach cells with Noxa antibody at 10 µg/mL.

Noxa Antibody - Background

Noxa Antibody: Apoptosis is related to many diseases and development. The p53 tumor-suppressor protein induces apoptosis through transcriptional activation of several genes including p53R2, p53AIP1, and PUMA. A new p53 target gene, Noxa, was recently identified, which encodes a protein belonging to the subfamily of BH3-only proapoptotic proteins. Noxa and PUMA are both transcriptional targets of p53 and BH3-only proteins. X-ray irradiation increased p53-dependent Noxa mRNA and protein levels. Noxa, when ectopically expressed, interacted with anti-apoptotic Bcl-2 family members, resulting in the activation of caspase-9. Noxa, like PUMA, localized to mitochondria and induces apoptosis in response to p53. Noxa and PUMA may represent direct mediators of p53-induced apoptosis. Increased levels of p53 and its target gene Noxa was found in the impaired tumor development.

Noxa Antibody - References

Oda E, Ohki R, Murasawa H, et al. Noxa, a BH3-only member of the Bcl-2 family and candidate mediator of p53-induced apoptosis. *Science* 2000; 288:1053-8.
Nakano K and Vousden KH. PUMA, a novel proapoptotic gene, is induced by p53. *Mol. Cell* 2001; 7:683-94.
Yu J, Zhang L, Hwang PM, et al. PUMA induces the rapid apoptosis of colorectal cancer cells. *Mol. Cell* 2001; 7:673-82.
Eferl R, Ricci R, Kenner L, et al. Liver tumor development. c-Jun antagonizes the proapoptotic activity of p53. *Cell* 2003; 112:181-92.