

PID Antibody
Catalog # ASC10138**Specification**

PID Antibody - Product Information

Application	WB, ICC, IF
Primary Accession	O94776
Other Accession	AAG02241 , 9931638
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 73 kDa

Application Notes	Observed: 75 kDa KDa PID antibody can be used for detection of PID by Western blot at 1 µg/mL. Antibody can also be used for immunocytochemistry starting at 10 µg/mL. For immunofluorescence start at 10 µg/mL.
-------------------	--

PID Antibody - Additional InformationGene ID **9219****Other Names**

PID Antibody: PID, MTA1L1, PID, Metastasis-associated protein MTA2, Metastasis-associated 1-like 1, MTA1-L1 protein, metastasis associated 1 family, member 2

Target/Specificity

MTA2; PID antibody is predicted to not cross-react with MTA2

Reconstitution & Storage

PID antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

PID Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PID Antibody - Protein Information**Name** MTA2**Synonyms** MTA1L1, PID**Function**

May function as a transcriptional coregulator (PubMed:16428440, PubMed:<a

<http://www.uniprot.org/citations/28977666> target="_blank">28977666). Acts as a component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin (PubMed:16428440, PubMed:28977666).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00512, ECO:0000255|PROSITE-ProRule:PRU00624, ECO:0000269|PubMed:28977666, ECO:0000269|PubMed:33283408}

Tissue Location


Widely expressed.

PID Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PID Antibody - Images



Western blot analysis of PID expression in HeLa whole cell lysates in the absence (A) or presence (B) of blocking peptide with PID antibody at 1 μ g/mL.



Immunocytochemistry staining of HeLa using PID antibody at 10 µg/mL.



Immunofluorescence of PID in HeLa cells with PID antibody at 10 µg/mL.

PID Antibody - Background

PID Antibody: The p53 tumor-suppressor gene integrates numerous signals that control cell life and death. Several novel molecules involved in p53 pathway, including Chk2, p53R2, p53AIP1, Noxa, PIDD, and PID/MTA2, were recently discovered. The transcriptional activity of p53 is modulated by protein stability and acetylation. PID/MTA2, also termed MTA1-L1, was found to be a subunit of nucleosome remodeling and deacetylating (NRD/NuRD) complex. PID/MTA2 modulates the enzymatic activity of the histone deacetylase complex and its expression reduces the levels of acetylated p53. Deacetylation of p53 by PID/MTA2 represses p53-dependent transcriptional activation and modulates p53-mediated cell growth arrest and apoptosis. PID/MTA2 is ubiquitously expressed in human tissues.

PID Antibody - References

Matsuoka S, Huang M, and Elledge SJ. Linkage of ATM to cell cycle regulation by the Chk2 protein kinase. *Science* 1998; 282:1893-7.
Tanaka H, Arakawa H, Yamaguchi T, et al. A ribonucleotide reductase gene involved in a p53-dependent cell-cycle checkpoint for DNA damage. *Nature* 2000; 404:42-9.
Oda E, Ohki R, Murasawa H, et al. Noxa, a BH3-only member of the Bcl-2 family and candidate mediator of p53-induced apoptosis. *Science* 2000; 288:1053-8.
Oda K, Arakawa H, Tanaka T, et al. p53AIP1, a potential mediator of p53-dependent apoptosis, and its regulation by Ser-46-phosphorylated p53. *Cell* 2000;102:849-62.