10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Mcl-1 Antibody
Catalog # ASC10305

Specification

Mcl-1 Antibody - Product Information

Application	**WB, ICC, IF**
Primary Accession	Q07820
Other Accession	NP_068779, 11386165
Reactivity	**Human, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**IgG**
Calculated MW	**Predicted: 22, 39 kDa**
	Observed: 39 kDa KDa
Application Notes	**Mcl-1 antibody can be used for detection of isoforms Mcl-1L and Mcl-1ES by Western blot at 1 to 2 µg/mL. Antibody can also be used for immunocytochemistry starting at 2 µg/mL. For immunofluorescence start at 10 µg/mL.**

Mcl-1 Antibody - Additional Information

Gene ID **4170**

Other Names
Mcl-1 Antibody: TM, EAT, MCL1L, MCL1S, Mcl-1, BCL2L3, MCL1-ES, bcl2-L-3, mcl1/EAT, Induced myeloid leukemia cell differentiation protein Mcl-1, Bcl-2-like protein 3, Bcl2-L-3, myeloid cell leukemia sequence 1 (BCL2-related)

Target/Specificity
MCL1; This Mcl-1 antibody detects isoforms Mcl-1L and Mcl-1ES.

Reconstitution & Storage
Mcl-1 antibody can be stored at 4℃ for three months and -20℃, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions
Mcl-1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Mcl-1 Antibody - Protein Information

Name MCL1

Synonyms BCL2L3

Function

Involved in the regulation of apoptosis versus cell survival, and in the maintenance of viability but not of proliferation. Mediates its effects by interactions with a number of other regulators of apoptosis. Isoform 1 inhibits apoptosis. Isoform 2 promotes apoptosis.

Cellular Location
Membrane; Single-pass membrane protein. Cytoplasm. Mitochondrion. Nucleus, nucleoplasm
Note=Cytoplasmic, associated with mitochondria

Mcl-1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Mcl-1 Antibody - Images



Western blot analysis of Mcl-1 in Raji cell lysate with Mcl-1 antibody at 0.5 µg/mL.



Immunocytochemistry staining of Raji cells using Mcl-1 antibody at 2 µg/mL.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Immunofluorescence of Mcl-1 in Raji cells with Mcl-1 antibody at 10 µg/mL.

Mcl-1 Antibody - Background

Mcl-1 Antibody: Myeloid cell leukimia-1 (Mcl-1) is a member of the Bcl-2 family of proteins that can act to promote cell survival. While the mechanism by which Mcl-1 inhibits apoptosis is not known, it is thought that it may heterodimerize and neutralize pro-apoptotic members of the Bcl-2 family such as Bim or Bak. Mcl-1 was originally identified in differentiating myeloid cells, but has since been shown to be expressed in multiple cell types. Mcl-1 is essential for embryogenesis and for the development and maintenance of B and T lymphocytes in animals. Mcl-1 exists as at least three distinct isoforms designated Mcl-1L, Mcl-1S and Mcl-1ES. In marked contrast to the larger isoform of Mcl-1, overexpression of Mcl-1S promotes cell death.

Mcl-1 Antibody - References

Edwards SW, Derouet M, Howse M, et al. Regulation of neutrophil apoptosis by Mcl-1. Biochem. Soc. Trans. 2004; 32:489-92.
Cuconati A, Mukherjee C, Perez D, et al. DNA damage response and MCL-1 destruction initiate apoptosis in adenovirus-infected cells. Genes and Dev. 2003; 17:2922-32.
Kozopas KM, Yang T, Buchan HL, et al. MCL1, a gene expressed in programmed myeloid cell differentiation, has sequence similarity to BCL2. Proc. Natl. Acad. Sci. USA 1993; 90:3516-20.