

IL-32 Antibody
Catalog # ASC10400

Specification

IL-32 Antibody - Product Information

Application	WB
Primary Accession	P24001
Other Accession	AAH09401 , 14424787
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	IL-32 antibody can be used for the detection of IL-32 by Western blot at 2.5 - 5 µg/mL.

IL-32 Antibody - Additional Information

Gene ID	9235
Other Names	IL-32 Antibody: NK4, TAIF, TAIFa, TAIFb, TAIFc, TAIFd, IL-32beta, IL-32alpha, IL-32delta, IL-32gamma, NK4, Interleukin-32, Natural killer cells protein 4, IL-32, interleukin 32

Target/Specificity

IL32; This antibody detects the largest isoform of IL-32.

Reconstitution & Storage

IL-32 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

IL-32 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

IL-32 Antibody - Protein Information

Name IL32

Synonyms NK4, TAIF

Function

Cytokine that may play a role in innate and adaptive immune responses. It induces various cytokines such as TNFA/TNF-alpha and IL8. It activates typical cytokine signal pathways of NF-kappa-B and p38 MAPK.

Cellular Location

Secreted.

Tissue Location


Selectively expressed in lymphocytes. Expression is more prominent in immune cells than in non-immune cells

IL-32 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IL-32 Antibody - Images



Western blot analysis of IL-32 in Jurkat cell lysate with IL-32 antibody at (A) 2.5 and (B) 5 μ g/mL.

IL-32 Antibody - Background

IL-32 Antibody: Interleukin-32 (IL-32) was initially identified as a transcript (NK4) that is selectively expressed in lymphocytes and NK cells and whose expression is increased following activation by IL-2. It was later re-isolated from an IL-18-treated lung carcinoma cell line and re-named IL-32. IL-32 is unusual in that it does not share sequence homology with known cytokine families and is highly expressed in immune tissues, existing in at least four differentially spliced isoforms. Because treatment of human monocytic and mouse macrophage cells with IL-32 induces several proinflammatory cytokines such as TNF- α , IL-8 and MIP-2, and because it is also induced in human peripheral lymphocyte cells after mitogen stimulation and in epithelial cells by IFN γ , it has been suggested that IL-32 may play a role in autoimmune and inflammatory diseases such as rheumatoid arthritis.

IL-32 Antibody - References

Dahl CA, Schall RP, He HL, et al. Identification of a novel gene expressed in activated natural killer cells and T cells. *J. Immunol.* 1992; 148:597-603.

Kim S-H, Han S-Y, Azam T, et al. Interleukin-32: a cytokine and inducer of TNF- α . *Immunity* 2005; 22:131-42.

Cagnard N, Letourneur F, Essabani A, et al. Interleukin-32, CCL2, PF4F1 and GFD10 are the only cytokine/chemokine genes differentially expressed by in vitro cultured rheumatoid and

osteoarthritis fibroblast-like synoviocytes. Eur. Cyto. Network 2005; 16:289-92.