

Hrk Antibody
Catalog # ASC10408**Specification**

Hrk Antibody - Product Information

Application	WB
Primary Accession	O00198
Other Accession	NP_003797 , 4504493
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 10 kDa

Application Notes	Observed: 12 kDa KDa Hrk antibody can be used for the detection of Hrk by Western blot at 2.5 - 5 µg/mL.
-------------------	---

Hrk Antibody - Additional InformationGene ID **8739****Other Names**

Hrk Antibody: DP5, HAKAKIRI, BID3, Activator of apoptosis harakiri, BH3-interacting domain-containing protein 3, harakiri, BCL2 interacting protein (contains only BH3 domain)

Target/Specificity

HRK;

Reconstitution & Storage

Hrk antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

Hrk Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Hrk Antibody - Protein Information**Name** HRK**Synonyms** BID3**Function**

Promotes apoptosis.

Cellular Location


Membrane; Single-pass membrane protein. Mitochondrion

Hrk Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Hrk Antibody - Images



Western blot analysis of Hrk in mouse pancreas tissue lysate with Hrk antibody at (A) 2.5 and (B) 5 µg/mL.

Hrk Antibody - Background

Hrk Antibody: Apoptosis plays a major role in normal organism development, tissue homeostasis, and removal of damaged cells. Hrk, a pro-apoptotic member of the Bcl-2 homology domain-3 (BH3)-only group of the Bcl-2 family of proteins, was also identified as novel protein induced during programmed neuronal death. It lacks significant homology to other Bcl-2 family members except for an 8-amino acid region that is similar to the BH3 motif of Bik. Hrk regulates apoptosis through interaction with the anti-apoptotic proteins Bcl-2 and Bcl-XL via this domain. It does not interact with the pro-apoptotic proteins Bax, Bak, or Bcl-XS. Hrk localizes to mitochondrial membranes in a pattern similar to that previously reported for Bcl-2 and Bcl-XL.

Hrk Antibody - References

Lockshin RA, Osborne B, and Zakeri Z. Cell death in the third millennium. *Cell Death Differ.* 2000; 7:2-7.

Imaizumi K, Tsuda M, Imai Y, et al. Molecular cloning of a novel polypeptide, DP5, induced during programmed neuronal death. *J. Biol. Chem.* 1997; 272:18842-8.

Inohara N, Ding L, Chen S, et al. harakiri, a novel regulator of cell death, encodes a protein that activates apoptosis and interacts selectively with survival-promoting proteins Bcl-2 and Bcl-XL. *EMBO J.* 1997; 16:1686-94.