

TRPC3 Antibody
Catalog # ASC10449**Specification**

TRPC3 Antibody - Product Information

Application	IF
Primary Accession	Q13507
Other Accession	CAA74083 , 7222
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 93 kDa

Application Notes

Observed: 100 kDa
TRPC3 antibody can be used for the detection of TRPC3 by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 10 µg/mL.

TRPC3 Antibody - Additional InformationGene ID **7222****Other Names**

TRPC3 Antibody: TRP3, TRP3, Short transient receptor potential channel 3, Transient receptor protein 3, TrpC3, transient receptor potential cation channel, subfamily C, member 3

Target/Specificity

TRPC3 antibody was raised against a 14 amino acid synthetic peptide from near the amino terminus of human TRPC3.

The immunogen is located within amino acids 120 - 170 of TRPC3.

Reconstitution & Storage

Antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.

Precautions

TRPC3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

TRPC3 Antibody - Protein Information**Name** TRPC3**Synonyms** TRP3**Function**

Forms a receptor-activated non-selective calcium permeant cation channel (PubMed:29726814, PubMed:30139744, PubMed:35051376, PubMed:9417057, PubMed:9930701, PubMed:10611319).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location


Expressed predominantly in brain and at much lower levels in ovary, colon, small intestine, lung, prostate, placenta and testis

TRPC3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRPC3 Antibody - Images



Immunofluorescence of BTK in Daudi cells with BTK antibody at 10 µg/mL.

TRPC3 Antibody - Background

TRPC3 Antibody: The mammalian transient receptor potential (TRP) superfamily can be divided into three major families including the "canonical TRP" (TRPC) family. The seven members of this family share the activation through PLC-coupled receptors and have been suggested to be components of receptor-regulated cation channels in different cell types. Furthermore, the members of the

TRPC3/6/7 subfamily can be activated by diacylglycerol analogs, suggesting a possible mechanism of activation of these channels by PLC-coupled receptors. TRPC3 encodes a Ca^{2+} -permeant channel that is agonist-activated but not store-operated or directly receptor-activated. TRPC3 physically interacts with TRPC6 and TRPC7 and forms functional tetrameric channels.

TRPC3 Antibody - References

Contell C, Birnbaumer V, Flockerzi V, et al. A unified nomenclature for the superfamily of TRP cation channels. *Mol. Cell* 2002; 9:229-31.

Trebak M, Vazquez G, Bird GSJ, et al. The TRPC3/6/7 subfamily of cation channels. *Cell Calcium* 2003; 33:451-61.

Zitt C, Obukhov AG, Strubing C, et al. Expression of TRPC3 in Chinese hamster ovary cells results in calcium-activated cation currents not related to store depletion. *J. Cell. Biol.* 1997; 1333-41.