

DRAM Antibody
Catalog # ASC10498**Specification**

DRAM Antibody - Product Information

Application	WB, IHC, IF
Primary Accession	Q8N682
Other Accession	AAH18435 , 22450862
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	DRAM antibody can be used for detection of DRAM by Western blot at 0.5 - 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 20 µg/mL.

DRAM Antibody - Additional Information

Gene ID 55332

Other Names

DRAM Antibody: DRAM, DRAM, DNA damage-regulated autophagy modulator protein 1, Damage-regulated autophagy modulator, damage-regulated autophagy modulator

Target/Specificity

DRAM;

Reconstitution & Storage

DRAM antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

DRAM Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

DRAM Antibody - Protein Information

Name DRAM1

Synonyms DRAM

Function

Lysosomal modulator of autophagy that plays a central role in p53/TP53-mediated apoptosis. Not involved in p73/TP73-mediated autophagy.

Cellular Location

Lysosome membrane; Multi-pass membrane protein

DRAM Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)


• [Flow Cytometry](#)

• [Cell Culture](#)


DRAM Antibody - Images



Western blot analysis of DRAM in 293 cell lysate with DRAM antibody at (A) 0.5 and (B) 1 µg/mL.



Immunohistochemistry of DRAM in human liver tissue with DRAM antibody at 2.5 µg/mL.



Immunofluorescence of DRAM in Human Liver cells with DRAM antibody at 20 µg/mL.

DRAM Antibody - Background

DRAM Antibody: Damage-regulated autophagy modulator (DRAM) is a p53 target gene encoding a lysosomal protein that induces autophagy, a process that degrades cytosolic proteins and organelles. It has been suggested that activation of DRAM by p53 is simultaneous to the activation by p53 of one or more proapoptotic genes such as PUMA, Bax, etc., and that the signaling pathways regulated by these genes promote a full cell death response. By itself, DRAM cannot induce apoptosis, but the fact that it is inactivated in certain cancers highlights the importance of DRAM and suggests that autophagy may play a more important role in cancer than initially suspected. At least two different isoforms of DRAM are known to exist.

DRAM Antibody - References

Crighton D, Wilkinson S, O'Prey J, et al. DRAM, a p53-induced modulator of autophagy, is critical for apoptosis. *Cell* 2006; 126:121-34.

Gozuacik D and Kimchi A. Autophagy as a cell death and tumor suppressor mechanism. *Oncogene*. 2004; 23:2891-906.

Crighton D, Wilkinson S, and Ryan KM. DRAM links autophagy to p53 and programmed cell death. *Autophagy* 2007; 3:72-4. (07-01D)