

PDL-2 Antibody
Catalog # ASC10510**Specification****PDL-2 Antibody - Product Information**

Application	WB
Primary Accession	O9BQ51
Other Accession	NP_079515 , 80380
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 30 kDa
Application Notes	Observed: 32 kDa KDa PD-L2 antibody can be used for detection of PD-L2 by Western blot at 0.5 - 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 20 µg/mL.

PDL-2 Antibody - Additional InformationGene ID **80380****Other Names**

PDL-2 Antibody: B7DC, Btdc, PDL2, CD273, PD-L2, PDCD1L2, bA574F11.2, B7DC, Programmed cell death 1 ligand 2, Butyrophilin B7-DC, PD-1 ligand 2, programmed cell death 1 ligand 2

Target/Specificity

PD-L2 antibody was raised against a 16 amino acid synthetic peptide from near the center of human PD-L2. The immunogen is located within amino acids 140 - 190 of PD-L2.

Reconstitution & Storage

PDL-2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

PDL-2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PDL-2 Antibody - Protein Information**Name** PDCD1LG2**Synonyms** B7DC, CD273, PDCD1L2, PDL2**Function**

Involved in the costimulatory signal, essential for T-cell proliferation and IFNG production in a PDCD1-independent manner. Interaction with PDCD1 inhibits T-cell proliferation by blocking cell cycle progression and cytokine production (By similarity).

Cellular Location

[Isoform 3]: Secreted [Isoform 1]: Cell membrane; Single-pass type I membrane protein {ECO:0000250|UniProtKB:Q9WUL5, ECO:0000305|PubMed:15340161}

Tissue Location


Highly expressed in heart, placenta, pancreas, lung and liver and weakly expressed in spleen, lymph nodes and thymus

PDL-2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PDL-2 Antibody - Images



Western blot analysis of UCMA in SW1353 cell lysate with UCMA antibody at 2.5 µg/mL.

PDL-2 Antibody - Background

PD-L2 Antibody: Cell-mediated immune responses are initiated by T lymphocytes that are themselves stimulated by cognate peptides bound to MHC molecules on antigen-presenting cells (APC). T-cell activation is generally self-limited as activated T cells express receptors such as PD-1 (also known as PDCD-1) that mediate inhibitory signals from the APC. PD-1 can bind two different but related ligands, PD-L1 and PD-L2, both of which are thought to act as a negative regulator of T cell activation. However, it has been suggested that PD-L2 can act to stimulate an immunogenic

response through and alternative receptor from PD-1.

PDL-2 Antibody - References

Holling TM, Schooten E, and van Den Elsing PJ. Function and regulation of MHC class II molecules in T-lymphocytes: of mice and men. *Hum. Immunol.* 2004; 65:282-90.

Ishida Y, Agata Y, Shibahara K, et al. Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death. *EMBO J.* 1992; 11:3887-95.

LaGier J and Pober JS. Immune accessory functions of human endothelial cells are modulated by overexpression of B7-H1 (PDL1). *Hum. Immunol.* 2006; 67:568-78.

Zhang Y, Chung Y, Bishop C, et al. Regulation of T cell activation and tolerance by PDL2. *Proc. Natl. Acad. Sci. USA* 2006; 103:11695-700.