

**OCC-1 Antibody**  
Catalog # ASC10833

**Specification**

---

**OCC-1 Antibody - Product Information**

Application	<b>WB, IHC</b>
Primary Accession	<a href="#">Q8TAD7</a>
Other Accession	<a href="#">AAH61920</a> , <a href="#">112180576</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>IgG</b>
Application Notes	<b>OCC-1 antibody can be used for detection of OCC-1 by Western blot at 2 - 4 µg/mL. In vitro transcription and translation of the OCC-1 cDNA yields polypeptides less than 9 kDa. Despite this, OCC-1 from cell lines is detected at ~28 kDa in SDS-PAGE. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL.</b>

**OCC-1 Antibody - Additional Information**

Gene ID	<b>387882</b>
<b>Target/Specificity</b>	
C12orf75;	

**Reconstitution & Storage**

OCC-1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

**Precautions**

OCC-1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**OCC-1 Antibody - Protein Information**

**Name** OCC1

**Synonyms** C12orf75

**Tissue Location**


High expression in placenta, skeletal muscle, kidney and pancreas tissues. Absent or very faint expression in heart, brain, lung and liver. Expressed during adipogenic differentiation of mesenchymal stem cells (at protein level)

## OCC-1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## OCC-1 Antibody - Images



Western blot analysis of OCC-1 in 293 cell lysate with OCC-1 antibody at (A) 2 µg/mL and (B) 4 µg/mL.



Immunohistochemistry of OCC-1 in human spleen tissue with OCC-1 antibody at 2.5 µg/mL.

## OCC-1 Antibody - Background

**OCC-1 Antibody:** OCC-1 (Overexpressed in colon carcinoma-1) was initially identified as a novel human cDNA overexpressed in a colon carcinoma cell line. Elevated levels of OCC-1 mRNA were observed in three of eight colon carcinomas, suggesting that OCC-1 may be a hallmark of only a subset of colon carcinomas. It is highly expressed in skeletal muscle, kidney, placenta, and pancreas tissues, with low to no expression in heart, brain, lung and liver tissues. At least two different OCC-1 mRNA species are known to exist.

**OCC-1 Antibody - References**

Pibouin L, Villaudy J, Ferbus D, et al. Cloning of the mRNA of overexpression in colon-carcinoma-1: a sequence overexpressed in a subset of colon carcinomas. *Cancer Genet. Cytogenet.* 2002; 133:55-60.