

RSPO1 Antibody
Catalog # ASC10908**Specification****RSPO1 Antibody - Product Information**

Application	WB
Primary Accession	Q2MKA7
Other Accession	NP_001033722 , 284654
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 20, 29 kDa

Application Notes	Observed: 26 kDa KDa RSPO1 antibody can be used for detection of RSPO1 by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 5 µg/mL.
-------------------	---

RSPO1 Antibody - Additional InformationGene ID **284654****Target/Specificity**

RSPO1 antibody was raised against a 16 amino acid synthetic peptide from near the amino terminus of human RSPO1. The immunogen is located within the first 50 amino acids of RSPO1.

Reconstitution & Storage

RSPO1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

RSPO1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

RSPO1 Antibody - Protein Information

Name RSPO1

Function

Activator of the canonical Wnt signaling pathway by acting as a ligand for LGR4-6 receptors (PubMed: [29769720](http://www.uniprot.org/citations/29769720)). Upon binding to LGR4-6 (LGR4, LGR5 or LGR6), LGR4-6 associate with phosphorylated LRP6 and frizzled receptors that are activated by extracellular Wnt receptors, triggering the canonical Wnt signaling pathway to increase expression of target genes. Also regulates the canonical Wnt/beta-catenin- dependent pathway and non-canonical Wnt signaling by acting as an inhibitor of

ZNRF3, an important regulator of the Wnt signaling pathway. Acts as a ligand for frizzled FZD8 and LRP6. May negatively regulate the TGF-beta pathway. Has a essential roles in ovary determination. Regulates Wnt signaling by antagonizing DKK1/KREM1- mediated internalization of LRP6 through an interaction with KREM1 (PubMed:17804805).

Cellular Location

Secreted. Nucleus {ECO:0000250|UniProtKB:Q9Z132} Note=Seems to mainly localize to nucleoli {ECO:0000250|UniProtKB:Q9Z132}

Tissue Location


Abundantly expressed in adrenal glands, ovary, testis, thyroid and trachea but not in bone marrow, spinal cord, stomach, leukocytes colon, small intestine, prostate, thymus and spleen.

RSPO1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RSPO1 Antibody - Images



Western blot analysis of XAF-1 in human brain tissue lysate with XAF-1 antibody at (A) 0.5 and (B) 1 µg/mL.

RSPO1 Antibody - Background

RSPO1 Antibody: RSPO1 is a member of a family of secreted growth factors that can operate through the canonical Wnt signaling pathway by stabilizing the intracellular beta-catenin, thereby regulating functions mediated by beta-catenin such as cell fate decisions and embryonic patterning.

RSPO1 was recently identified through linkage analysis to be involved in sex determination and mammalian ovarian development. RSPO1 is thought to regulate cellular responsiveness to Wnt ligands by modulating the cell-surface expression of the Wnt co-receptor LRP6 by interfering with the DKK/Kremen-mediated internalization of LRP6 through an interaction with Kremen, resulting in increased LRP6 cell-surface levels.

RSPO1 Antibody - References

Kamata T, Katsube K, Michikawa M, et al. R-spondin, a novel gene with thrombospondin type I domain, was expressed in the dorsal neural tube and affected Wnts mutants. *Biochim. Biophys. Acta* 2004; 1676:51-61.

Kim KA, Zhao J, Andarmani S, et al. R-spondin proteins: a novel link to beta-catenin signaling. *Cell Cycle* 2006; 5:23-6.

Parma P, Radi O, Vidal V, et al. R-spondin1 is essential in sex determination, skin differentiation and malignancy. *Nat. Genet.* 2006; 38:1304-9.

Chassot AA, Ranc F, Gregoire EP, et al. Activation of b-catenin signalling by Rspo1 controls differentiation of the mammalian ovary. *Hum. Mol. Genet.* 2008; 17:1278-91.