

# **SYNGR1 Antibody**

Catalog # ASC11046

## **Specification**

## **SYNGR1** Antibody - Product Information

Application
Primary Accession
Other Accession
Reactivity
Host
Clonality
Isotype
Application Notes

WB, IHC, IF 043759 CAA05322, 2959866

Human, Mouse, Rat Rabbit

Polyclonal

IgG

SYNGR1 antibody can be used for

detection of SYNGR1 by Western blot at 1 μg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 μg/mL. For immunofluorescence start at 20

μg/mL.

## **SYNGR1** Antibody - Additional Information

Gene ID 9145

Target/Specificity

SYNGR1:

#### **Reconstitution & Storage**

Antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.

#### **Precautions**

SYNGR1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

#### SYNGR1 Antibody - Protein Information

### Name SYNGR1 (HGNC:11498)

#### **Function**

May play a role in regulated exocytosis. Modulates the localization of synaptophysin/SYP into synaptic-like microvesicles and may therefore play a role in synaptic-like microvesicle formation and/or maturation (By similarity). Involved in the regulation of short- term and long-term synaptic plasticity (By similarity).

#### **Cellular Location**

Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane {ECO:0000250|UniProtKB:Q62876}; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q62876}. Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV




# **SYNGR1 Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# **SYNGR1 Antibody - Images**




Western blot analysis of SYNGR1 in rat brain tissue lysate with SYNGR1 antibody at 1  $\mu$ g/mL in (A) the absence and (B) the presence of blocking peptide.



Immunohistochemistry of SYNGR1 in rat brain tissue with SYNGR1 antibody at 2.5 μg/mL.





Immunofluorescence of SYNGR1 in rat brain tissue with SYNGR1 antibody at 20 µg/mL.

### SYNGR1 Antibody - Background

SYNGR1 Antibody: Synaptogyrins comprise a family of tyrosine-phosphorylated membrane proteins with two neuronal (SYNGR1 and SYNGR3) and one ubiquitous (SYNGR2) members. SYNGR1 and -3 are synaptic vesicle proteins, residing in some cases on the same synaptic vesicle, and are thought to be involved in the regulation of neurotransmitter release. SYNGR2, by contrast, is absent from synaptic vesicles. The role and localization of a fourth synaptogyrin, SYNGR4, is unclear. The gene for SYNGR1is located at chromosome 22q13, a region linked to schizophrenia; however, there is mixed evidence suggesting that mutations in SYNGR1 might be associated with schizophrenia.

# **SYNGR1 Antibody - References**

Kedra D, Pan HQ, Seroussi E, et al. Characterization of the human synaptogyrin gene family. Hum. Genet.1998; 103:131-41.

Egaña LA, Cuevas RA, Baust TB, et al. Physical and functional interaction between the dopamine transporter and the synaptic vesicle protein synaptogyrin-3. J. Neurosci.2009; 29:4592-604. Cheng MC and Chen CH. Identification of rare mutations of synaptogyrin 1 gene in patients with schizophrenia. J. Psychiatr. Res.2007; 41:1027-31.

Wang Y, Yu L, Zhao T, et al. No association between bipolar disorder and syngr1 or synapsin II polymorphisms in the Han Chinese population. Psychiatry Res.2009; 169:167-8.