

**CDX2 Antibody**  
Catalog # ASC11062**Specification****CDX2 Antibody - Product Information**

|                   |   |
|-------------------|---|
| Application       | WB, IHC, IF   |
| Primary Accession | <a href="#">Q99626</a>  |
| Other Accession   | <a href="#">CAA74038</a> , <a href="#">2193875</a>  |
| Reactivity        | Human, Mouse, Rat   |
| Host              | Rabbit  |
| Clonality         | Polyclonal  |
| Isotype           | IgG   |
| Application Notes | CDX2 antibody can be used for detection of CDX2 by Western blot at 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL. |

**CDX2 Antibody - Additional Information**

|                    |       |
|--------------------|-------|
| Gene ID            | 1045  |
| Target/Specificity | CDX2; |

**Reconstitution & Storage**

Antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.

**Precautions**

CDX2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**CDX2 Antibody - Protein Information**

**Name** CDX2

**Synonyms** CDX3

**Function**

Transcription factor which regulates the transcription of multiple genes expressed in the intestinal epithelium (By similarity). Binds to the promoter of the intestinal sucrase-isomaltase SI and activates SI transcription (By similarity). Binds to the DNA sequence 5'-ATAAAACTTAT-3' in the promoter region of VDR and activates VDR transcription (By similarity). Binds to and activates transcription of LPH (By similarity). Activates transcription of CLDN2 and intestinal mucin MUC2 (By similarity). Binds to the 5'-AATTTTTTACAACACCT-3' DNA sequence in the promoter region of CA1 and activates CA1 transcription (By similarity). Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestine. Binds preferentially to methylated DNA (PubMed:<a href="http://www.uniprot.org/citations/28473536" target="\_blank">28473536</a>).

### Cellular Location

Nucleus {ECO:0000250|UniProtKB:P43241}.

### Tissue Location


Detected in small intestine, colon and pancreas.

### CDX2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### CDX2 Antibody - Images



Western blot analysis of CDX2 in mouse brain tissue lysate with CDX2 antibody at 1 µg/mL.



Immunohistochemistry of CDX2 in rat brain tissue with CDX2 antibody at 5 µg/mL.



Immunofluorescence of CDX2 in rat brain tissue with CDX2 antibody at 20 µg/mL.

### **CDX2 Antibody - Background**

CDX2 Antibody: CDX2 is a member of the caudal homeobox family of transcription factors that also includes the proteins CDX1 and CDX4. CDX2 is strongly expressed in the trophectoderm in the blastocyst with expression maintained within the proliferating extra-embryonic ectoderm during development. It is required for the repression of the transcription of the POU5F1/Oct4 and NANOG genes in the outer cells of the blastocyst and is thus essential for the segregation of the inner cell mass and trophectoderm at the blastocyst stage. Aberrant expression of CDX2 is associated with numerous types of carcinomas, including intestinal metaplasia, gastric carcinoma, and acute myeloid leukemia due to deregulated HOX gene expression.

### **CDX2 Antibody - References**

Beck F, Erler T, Russell A, et al. Expression of CDX-2 in the mouse embryo and placenta: possible role in patterning of the extra-embryonic membranes. *Dev. Dyn.*1995; 204:219-27.  
Strumpf D, Mao CA, Yamanaka Y, et al. Cdx2 is required for correct cell fate specification and differentiation of trophectoderm in the mouse blastocyst. *Dev.*2005; 132:2093-102.  
Ishikawa A, Sasaki M, Ohira S, et al. Aberrant expression of CDX2 is closely related to the intestinal metaplasia and MUC2 expression in intraductal papillary neoplasm of the liver in hepatolithiasis. *Lab. Invest.*2004; 84:629-38.  
Silberg DG, Sullivan J, Kang E, et al. Cdx2 ectopic expression induces gastric intestinal metaplasia in transgenic mice. *Gastroenterology*2002; 122:689-96.