

SPT1 Antibody
Catalog # ASC11322**Specification****SPT1 Antibody - Product Information**

Application	IF
Primary Accession	O15269
Other Accession	NP_6406 , 10558
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	SPT1 antibody can be used for detection of SPT1 by Western blot at 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL.

SPT1 Antibody - Additional InformationGene ID **10558****Target/Specificity**

SPT1 antibody was raised against a 17 amino acid synthetic peptide near the carboxy terminus of human SPT1. The immunogen is located within amino acids 380 - 430 of SPT1.

Reconstitution & Storage

SPT1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

SPT1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

SPT1 Antibody - Protein Information

Name SPTLC1

Synonyms LCB1

Function

Component of the serine palmitoyltransferase multisubunit enzyme (SPT) that catalyzes the initial and rate-limiting step in sphingolipid biosynthesis by condensing L-serine and activated acyl-CoA (most commonly palmitoyl-CoA) to form long-chain bases. The SPT complex is also composed of SPTLC2 or SPTLC3 and SPTSSA or SPTSSB. Within this complex, the heterodimer with SPTLC2 or SPTLC3 forms the catalytic core (PubMed: [19416851](http://www.uniprot.org/citations/19416851), PubMed: [33558762](http://www.uniprot.org/citations/33558762), PubMed: [36170811](http://www.uniprot.org/citations/36170811)). The composition of the serine palmitoyltransferase (SPT) complex determines the substrate preference (PubMed:

[19416851](http://www.uniprot.org/citations/19416851), PubMed:<[33558762](http://www.uniprot.org/citations/33558762)>). The SPTLC1-SPTLC2-SPTSSA complex shows a strong preference for C16-CoA substrate, while the SPTLC1-SPTLC3-SPTSSA isozyme uses both C14-CoA and C16-CoA as substrates, with a slight preference for C14-CoA (PubMed:<[19416851](http://www.uniprot.org/citations/19416851)>, PubMed:<[19648650](http://www.uniprot.org/citations/19648650)>). The SPTLC1-SPTLC2-SPTSSB complex shows a strong preference for C18-CoA substrate, while the SPTLC1-SPTLC3-SPTSSB isozyme displays an ability to use a broader range of acyl-CoAs, without apparent preference (PubMed:<[19416851](http://www.uniprot.org/citations/19416851)>, PubMed:<[19648650](http://www.uniprot.org/citations/19648650)>, PubMed:<[33558761](http://www.uniprot.org/citations/33558761)>, PubMed:<[33558762](http://www.uniprot.org/citations/33558762)>). Required for adipocyte cell viability and metabolic homeostasis (By similarity).

Cellular Location

Endoplasmic reticulum membrane; Single-pass membrane protein
{ECO:0000250|UniProtKB:O35704}

Tissue Location


Widely expressed. Not detected in small intestine.

SPT1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SPT1 Antibody - Images



Immunofluorescence of KIR2DS2 in 293 cells with KIR2DS2 antibody at 20 µg/ml.

SPT1 Antibody - Background

SPT1 Antibody: Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid biosynthesis. It converts L-serine and palmitoyl-CoA to 3-oxosphinganine with pyridoxal 5'-phosphate as a cofactor. SPT1 is the long chain base subunit 1 of mammalian serine palmitoyltransferase. SPT1 is not catalytically active but is necessary for the stabilization of the SPT2 subunit and anchoring the holoenzyme to the cytosolic face of the endoplasmic reticulum. Missense mutations in this gene have been identified in patients with hereditary sensory neuropathy type 1 (HSAN1). These mutations induce a shift in the substrate specificity of the holoenzyme, leading to the formation and accumulation of two neurotoxic sphingolipids.

SPT1 Antibody - References

- Batheja AD, Uhlinger DJ, Carton JM, et al. Characterization of serine palmitoyltransferase in normal human tissues. *J. Histochem. & Cytochem.* 2003; 51:687-96.
- Hanada K, Hara T, Nishijima N, et al. A mammalian homolog of the yeast LCB1 encodes a component of serine palmitoyltransferase, the enzyme catalyzing the first step in sphingolipid synthesis. *J. Biol. Chem.* 1997; 272:32108-14.
- Yasuda S, Nishijima N, and Hanada K. Localization, topology, and function of the LCB1 subunit of serine palmitoyltransferase in mammalian cells. *J. Biol. Chem.* 2003; 278:4176-83.
- Penno A, Reilly MM, Houlden H, et al. Hereditary sensory neuropathy type 1 is caused by the accumulation of two neurotoxic sphingolipids. *J. Biol. Chem.* 2010; 285:11178-87.