

BMP15 Antibody
Catalog # ASC11499**Specification**

BMP15 Antibody - Product Information

Application	WB, ICC, IF
Primary Accession	O95972
Other Accession	NP_005439 , 257743454
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	BMP15 antibody can be used for detection of BMP15 by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunocytochemistry starting at 2.5 µg/mL. For immunofluorescence start at 2.5 µg/mL.

BMP15 Antibody - Additional Information

Gene ID 9210

Target/Specificity

BMP15; It is closely related to GDF9.

Reconstitution & Storage

BMP15 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

BMP15 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

BMP15 Antibody - Protein Information

Name BMP15

Synonyms GDF9B

Function

May be involved in follicular development. Oocyte-specific growth/differentiation factor that stimulates folliculogenesis and granulosa cell (GC) growth.

Cellular Location


Secreted.

BMP15 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


BMP15 Antibody - Images



Western blot analysis of BMP15 in 3T3 cell lysate with BMP15 antibody at (A) 1 and (B) 2 µg/ml



Immunocytochemistry of BMP15 in 3T3 cells with BMP15 antibody at 2.5 µg/mL.



BMP15 Antibody - Background

BMP15 Antibody: BMP15, or GDF9B is a member of the bone morphogenetic protein family which is part of the transforming growth factor-beta superfamily that are involved in embryonic development and adult tissue homeostasis. BMP15 is expressed exclusively in the oocyte. It is an oocyte-specific growth/differentiation factor that may be involved in oocyte maturation and follicular development. Defects in BMP15 are the cause of ovarian dysgenesis type 2. BMP15 could be used as an oogenesis marker to track human amniotic fluid stem cells differentiation into the oocyte-like cells.

BMP15 Antibody - References

- Massagué J. The transforming growth factor-beta family. *Ann. Rev. Cell. Biol.* 1990; 6:597-641
- Aaltonen J, Laitinen MP, Vuojolainen K, et al. Human growth differentiation factor 9 (GDF-9) and its novel homolog GDF-9B are expressed in oocytes during early folliculogenesis. *J. Clin. Endocrinol. Metab.* 1999; 84:2744-50.
- Laitinen M, Vuojolainen K, Jaatinen R, et al. A novel growth differentiation factor-9 (GDF-9) related factor is coexpressed with GDF-9 in mouse oocytes during folliculogenesis. *Mech. Dev.* 1998; 78:135-40.
- Galloway SM, Gregan SM, Wilson T et al. Bmp15 mutations and ovarian function. *Mol. Cell Endocrinol.* 2002; 191:15-8.