

**TAF9 Antibody**  
Catalog # ASC11772**Specification**

---

**TAF9 Antibody - Product Information**

|                   |   |
|-------------------|---|
| Application       | WB, IHC, IF   |
| Primary Accession | <a href="#">Q16594</a>                              |
| Other Accession   | <a href="#">NP_003178</a> , <a href="#">4507351</a> |
| Reactivity        | Human, Mouse, Rat                                   |
| Host              | Rabbit  |
| Clonality         | Polyclonal  |
| Isotype           | IgG   |
| Calculated MW     | Predicted: 29 kDa                                   |

|                   |   |
|-------------------|---|
| Application Notes | <b>Observed: 28 kDa KDa</b><br>TAF9 antibody can be used for detection of TAF9 by Western blot at 1 - 2 µg/ml. Antibody can also be used for Immunohistochemistry at 5 µg/mL. For Immunofluorescence start at 20 µg/mL. |
|-------------------|---|

**TAF9 Antibody - Additional Information**Gene ID **6880****Target/Specificity**

TAF9; TAF9 antibody is human, mouse and rat reactive. Multiple isoforms of TAF9 are known to exist. TAF9 is predicted to not cross-react with other TAF family members.

**Reconstitution & Storage**

TAF9 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

**Precautions**

TAF9 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**TAF9 Antibody - Protein Information**

Name TAF9

Synonyms TAF2G, TAFII31

**Function**

The TFIID basal transcription factor complex plays a major role in the initiation of RNA polymerase II (Pol II)-dependent transcription (PubMed: [33795473](http://www.uniprot.org/citations/33795473)). TFIID recognizes and binds promoters with or without a TATA box via its subunit TBP, a TATA-box-binding protein, and promotes assembly of the pre-initiation complex (PIC) (PubMed: [33795473](http://www.uniprot.org/citations/33795473)). The TFIID complex consists of TBP and TBP-associated factors (TAFs), including TAF1, TAF2, TAF3, TAF4, TAF5, TAF6, TAF7, TAF8, TAF9, TAF10, TAF11, TAF12 and

TAF13 (PubMed:<a href="http://www.uniprot.org/citations/33795473" target="\_blank">33795473</a>). TAF9 is also a component of the TBP-free TAFII complex (TFTC), the PCAF histone acetylase complex and the STAGA transcription coactivator-HAT complex (PubMed:<a href="http://www.uniprot.org/citations/15899866" target="\_blank">15899866</a>). TAF9 and its paralog TAF9B are involved in transcriptional activation as well as repression of distinct but overlapping sets of genes (PubMed:<a href="http://www.uniprot.org/citations/15899866" target="\_blank">15899866</a>). Essential for cell viability (PubMed:<a href="http://www.uniprot.org/citations/15899866" target="\_blank">15899866</a>). May have a role in gene regulation associated with apoptosis (PubMed:<a href="http://www.uniprot.org/citations/15899866" target="\_blank">15899866</a>).

### Cellular Location


Nucleus

### TAF9 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### TAF9 Antibody - Images



Western blot analysis of TAF9 in HepG2 cell lysate with TAF9 antibody at (A) 1 and (B) 2 µg/ml.



Immunohistochemistry of TAF9 in mouse liver tissue with TAF9 antibody at 5 µg/mL.



Immunofluorescence of TAF9 in mouse liver tissue with TAF9 antibody at 20 µg/mL.

### **TAF9 Antibody - Background**

Initiation of transcription by RNA polymerase II is coordinated by the transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and the TBP-associated factors (TAFs). TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation (1). TAF9 is one of the smaller subunits of TFIID that binds to the basal transcription factor GTF2B as well as to several transcriptional activators such as p53 and VP16 (2).

### **TAF9 Antibody - References**

Zawel L and Reinberg D. Advances in RNA polymerase II transcription. *Curr. Opin. Cell Biol.* 1992; 4:488-95.  
Klemm RD, Goodrich JA, Zhou S, et al. Molecular cloning and expression of the 32-kDa subunit of human TFIID reveals interactions with VP16 and TFIIB that mediate transcriptional activation. *Proc. Natl. Acad. Sci. USA* 1995; 92:5788-92.