

**Vinculin Antibody**  
Catalog # ASC11803**Specification****Vinculin Antibody - Product Information**

Application	IF, IHC
Primary Accession	<a href="#">P18206</a>
Other Accession	<a href="#">NP_003364</a> , <a href="#">7414</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 120 kDa
Application Notes	Observed: 117 kDa KDa Vinculin antibody can be used for detection of Vinculin by Western blot at 0.5 - 1 µg/ml. For immunofluorescence start at 20 µg/mL.

**Vinculin Antibody - Additional Information**Gene ID **7414****Target/Specificity**

Vinculin antibody was raised against a 23 amino acid peptide near the center of human Vinculin. The immunogen is located within amino acids 820 - 870 of Vinculin.

**Reconstitution & Storage**

Vinculin antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

**Precautions**

Vinculin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Vinculin Antibody - Protein Information**

Name VCL

**Function**

Actin filament (F-actin)-binding protein involved in cell- matrix adhesion and cell-cell adhesion. Regulates cell-surface E- cadherin expression and potentiates mechanosensing by the E-cadherin complex. May also play important roles in cell morphology and locomotion.

**Cellular Location**

Cell membrane {ECO:0000250|UniProtKB:P12003}; Peripheral membrane protein {ECO:0000250|UniProtKB:P12003}; Cytoplasmic side {ECO:0000250|UniProtKB:P12003}. Cell junction, adherens junction {ECO:0000250|UniProtKB:P12003}. Cell junction, focal adhesion {ECO:0000250|UniProtKB:P12003}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P85972}. Cell membrane, sarcolemma {ECO:0000250|UniProtKB:Q64727}; Peripheral membrane protein

{ECO:0000250|UniProtKB:Q64727}; Cytoplasmic side {ECO:0000250|UniProtKB:Q64727}. Cell projection, podosome {ECO:0000250|UniProtKB:Q64727}. Note=Recruitment to cell-cell junctions occurs in a myosin II-dependent manner. Interaction with CTNNB1 is necessary for its localization to the cell-cell junctions {ECO:0000250|UniProtKB:P12003}

#### Tissue Location


Metavinculin is muscle-specific.

#### Vinculin Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Vinculin Antibody - Images



Immunofluorescence of RAIDD in HeLa cells with RAIDD antibody at 5 µg/ml.



### **Vinculin Antibody - Background**

Vinculin is a cytoskeletal protein that plays an important role in the regulation of focal adhesions and embryonic development (1). Three structural vinculin domains include an amino-terminal head, a short flexible proline-rich region and a carboxy-terminal tail (2). Expression of vinculin were shown to be affected by the level of actin expression (2,3). Vinculin deficiencies are associated with a decrease in cell adhesion and an increase in cell motility, suggesting a possible role in metastatic growth (4). Defects in VCL are the cause of cardiomyopathy dilated type 1W (CMD1W) (5).

### **Vinculin Antibody - References**

- Burridge K, Fath K, Kelly T, et al. Focal adhesions: transmembrane junctions between the extracellular matrix and the cytoskeleton. *Annu. Rev. Cell Biol.* 1988; 4:487-525.
- Gilmore AP, Jackson P, Waites GT, et al. Further characterization of the talin-binding site in the cytoskeletal protein vinculin. *J. Cell Sci.* 1992; 103:719-31.
- Deakin NO, Ballestrem C, and Turner CE. Paxillin and Hic-5 interaction with vinculin is differentially regulated by Rac1 and RhoA. *PLoS One* 2012; 7:e37990.
- Goldmann WH, Auernheimer V, Thievensen I, et al. Vinculin, cell mechanics and tumour cell invasion. *Cell Biol. Int.* 2013; Feb 1.