

**ARID1A Antibody**  
Catalog # ASC11922**Specification****ARID1A Antibody - Product Information**

|                   |   |
|-------------------|---|
| Application       | IHC   |
| Primary Accession | <a href="#">O14497</a>  |
| Other Accession   | <a href="#">NP_006006</a> , <a href="#">8289</a>  |
| Reactivity        | Human, Mouse, Rat   |
| Host              | Rabbit  |
| Clonality         | Polyclonal  |
| Isotype           | IgG   |
| Calculated MW     | Predicted: 133, 227, 251 kDa; Observed: 130, 260 kDa  |
| Application Notes | ARID1A antibody can be used for detection of ARID1A by Western blot at 1 - 2 µg/mL. For immunofluorescence start at 20 µg/mL. |

**ARID1A Antibody - Additional Information**Gene ID **8289****Target/Specificity**

ARID1A antibody was raised against an 18 amino acid peptide near the amino terminus of human ARID1A. The immunogen is located within amino acids 580 - 630 of ARID1A.

**Reconstitution & Storage**

ARID1A antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

**Precautions**

ARID1A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**ARID1A Antibody - Protein Information**

Name ARID1A

Synonyms BAF250, BAF250A, C1orf4, OSA1, SMARCF1

**Function**

Involved in transcriptional activation and repression of select genes by chromatin remodeling (alteration of DNA-nucleosome topology). Component of SWI/SNF chromatin remodeling complexes that carry out key enzymatic activities, changing chromatin structure by altering DNA-histone contacts within a nucleosome in an ATP-dependent manner. Binds DNA non-specifically. Belongs to the neural progenitors- specific chromatin remodeling complex (npBAF complex) and the neuron- specific chromatin remodeling complex (nBAF complex). During neural development a switch from a stem/progenitor to a postmitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to postmitotic neurons requires a switch in subunit

composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth (By similarity).

#### Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00355, ECO:0000269|PubMed:11318604, ECO:0000269|PubMed:26614907}

#### Tissue Location


Highly expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, and PBL, and at a much lower level in heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas.

### ARID1A Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### ARID1A Antibody - Images



Immunohistochemistry of Neurturin in mouse brain tissue with Neurturin Antibody at 5 µg/mL.

### ARID1A Antibody - Background

The ARID1A protein is a member of the SWI/SNF family, whose members are thought to regulate transcription of certain genes by altering the chromatin structure around those genes. ARID1A is part of the large ATP-dependent chromatin remodeling complex SNF/SWI, which is required for transcriptional activation of genes normally repressed by chromatin (1). It possesses a DNA-binding

domain that can specifically bind an AT-rich DNA sequence known to be recognized by a SNF/SWI complex at the beta-globin locus. The C-terminus of the protein can stimulate glucocorticoid receptor-dependent transcriptional activation. It is thought that ARID1A confers specificity to the SNF/SWI complex and may recruit the complex to its targets through either protein-DNA or protein-protein interactions (2).

#### **ARID1A Antibody - References**

Martens JA and Winston F. Recent advances in understanding chromatin remodeling by Swi/Snf complexes. *Curr. Opin. Genet. Dev.* 2003; 13:136-42.

Nie Z, Xue Y, Yang D, et al. A specificity and targeting subunit of a human SWI/SNF family-related chromatin-remodeling complex. *Mol. Cell. Biol.* 2000; 20:8879-88.