

**Hexokinase 1 Antibody**  
Catalog # ASC11938**Specification****Hexokinase 1 Antibody - Product Information**

Application	WB, IHC, IF
Primary Accession	<a href="#">P19367</a>
Other Accession	<a href="#">NP_000179</a> , <a href="#">188497754</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 101 kDa; Observed: 103 kDa KDa
Application Notes	Hexokinase 1 antibody can be used for detection of Hexokinase 1 by Western blot at 1 - 2 µg/ml. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.

**Hexokinase 1 Antibody - Additional Information**Gene ID **3098****Target/Specificity**

HK1; Hexokinase 1 antibody is human, mouse and rat reactive. Multiple isoforms of Hexokinase 1 are known to exist.

**Reconstitution & Storage**

Hexokinase 1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

**Precautions**

Hexokinase 1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Hexokinase 1 Antibody - Protein Information**Name HK1 ([HGNC:4922](#))**Function**

Catalyzes the phosphorylation of various hexoses, such as D- glucose, D-glucosamine, D-fructose, D-mannose and 2-deoxy-D-glucose, to hexose 6-phosphate (D-glucose 6-phosphate, D-glucosamine 6-phosphate, D-fructose 6-phosphate, D-mannose 6-phosphate and 2-deoxy-D-glucose 6- phosphate, respectively) (PubMed:[1637300](http://www.uniprot.org/citations/1637300), PubMed:[25316723](http://www.uniprot.org/citations/25316723), PubMed:[27374331](http://www.uniprot.org/citations/27374331)). Does not phosphorylate N-acetyl-D-glucosamine (PubMed:

<http://www.uniprot.org/citations/27374331>). Mediates the initial step of glycolysis by catalyzing phosphorylation of D-glucose to D-glucose 6-phosphate (By similarity). Involved in innate immunity and inflammation by acting as a pattern recognition receptor for bacterial peptidoglycan (PubMed:[27374331](http://www.uniprot.org/citations/27374331)). When released in the cytosol, N-acetyl-D-glucosamine component of bacterial peptidoglycan inhibits the hexokinase activity of HK1 and causes its dissociation from mitochondrial outer membrane, thereby activating the NLRP3 inflammasome (PubMed:[27374331](http://www.uniprot.org/citations/27374331)).

#### Cellular Location

Mitochondrion outer membrane; Peripheral membrane protein. Cytoplasm, cytosol. Note=The mitochondrial-binding peptide (MBP) region promotes association with the mitochondrial outer membrane (Probable). Dissociates from the mitochondrial outer membrane following inhibition by N-acetyl-D-glucosamine, leading to relocation to the cytosol (PubMed:27374331).

#### Tissue Location


Isoform 2: Erythrocyte specific (Ref.6). Isoform 3: Testis-specific (PubMed:10978502). Isoform 4: Testis-specific (PubMed:10978502). {ECO:0000269|PubMed:10978502, ECO:0000269|Ref.6}

### Hexokinase 1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### Hexokinase 1 Antibody - Images



Western blot analysis of Hexokinase 1 in rat bladder tissue lysate with Hexokinase 1 antibody at (A) 1 and (B) 2 µg/ml.



Immunohistochemistry of Hexokinase 1 in rat stomach tissue with Hexokinase 1 antibody at 5 µg/mL.



Immunofluorescence of Hexokinase 1 in rat stomach tissue with Hexokinase 1 antibody at 20 µg/mL.

### Hexokinase 1 Antibody - Background

There are four major glucose-phosphorylating isoenzymes, designated Hexokinase 1 I, II, III, and IV (1). Hexokinase 1 activity is involved in the first step in several metabolic pathways including phosphorylation of glucose to produce glucose-6-phosphate, thus committing glucose to the glycolytic pathway (1,2). Hexokinase 1 2 is the predominant Hexokinase 1 isozyme expressed in insulin-responsive tissues such as skeletal muscle and its expression is insulin-responsive (3). It is involved in the increased rate of glycolysis seen in rapidly growing cancer cells (4).

### Hexokinase 1 Antibody - References

- Chen T, Ning D, Sun H, et al. Sequence Analysis and Molecular Characterization of Clonorchis sinensis Hexokinase 1, an Unusual Trimeric 50-kDa Glucose-6-Phosphate-Sensitive Allosteric Enzyme. *PLoS One* 2014; 9:e107940.
- Halestrap AP, Pereira GC, and Pasdois P. The role of Hexokinase 1 in cardioprotection-mechanism and potential for translation. *Br. J. Pharmacol.* 2014; epub.
- Roberts DJ and Miyamoto S. Hexokinase 1 II integrates energy metabolism and cellular protection: Akt acting on mitochondria and TORCing to autophagy. *Cell Death Differ.* 2014; epub.
- Wang L, Xiong H, Wu F, et al. Hexokinase 1 2-Mediated Warburg Effect Is Required for PTEN- and p53-Deficiency-Driven Prostate Cancer Growth. *Cell Rep.* 2014; 8:1461-74.