

ICOS Antibody
Catalog # ASC12119**Specification**

ICOS Antibody - Product Information

Application	WB, IHC, IF, E
Primary Accession	O9Y6W8
Other Accession	O9Y6W8
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG

ICOS Antibody - Additional Information

Gene ID	29851
Alias Symbol	ICOS

Other Names

ICOS Antibody: Inducible T Cell Costimulator, Activation-Inducible Lymphocyte Immunomediatory Molecule, Inducible T-Cell Costimulator, ALLIM, Inducible T-Cell Co-Stimulator, Inducible Costimulator, CD278 Antigen, CD278, CVID1

Reconstitution & Storage

ICOS antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

ICOS Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

ICOS Antibody - Protein Information

Name ICOS

Synonyms ALLIM

Function

Enhances all basic T-cell responses to a foreign antigen, namely proliferation, secretion of lymphokines, up-regulation of molecules that mediate cell-cell interaction, and effective help for antibody secretion by B-cells. Essential both for efficient interaction between T and B-cells and for normal antibody responses to T-cell dependent antigens. Does not up-regulate the production of interleukin- 2, but superinduces the synthesis of interleukin-10. Prevents the apoptosis of pre-activated T-cells. Plays a critical role in CD40- mediated class switching of immunoglobulin isotypes (By similarity).

Cellular Location

[Isoform 1]: Cell membrane; Single- pass type I membrane protein

Tissue Location

Activated T-cells. Highly expressed on tonsillar T- cells, which are closely associated with B-cells in the apical light zone of germinal centers, the site of terminal B-cell maturation Expressed at lower levels in thymus, lung, lymph node and peripheral blood leukocytes. Expressed in the medulla of fetal and newborn thymus

ICOS Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ICOS Antibody - Images