

WNT3 Antibody
Rabbit Anti-Human WNT3 Polyclonal
Catalog # ASM10632

Specification

WNT3 Antibody - Product Information

Application	IHC
Primary Accession	P56703
Other Accession	NP_110380.1
Host	Rabbit
Clonality	Polyclonal
Target/Specificity	
WNT3	

Other Names

WNT 3 proto oncogene protein precursor Antibody, wingless type MMTV integration site family member 3 Antibody, Wnt 3 proto oncogene protein Antibody, MGC138321 Antibody, WNT3_HUMAN Antibody, Proto-oncogene Int-4 homolog Antibody, MGC138323 Antibody, MGC131950 Antibody, INT4 Antibody, Proto-oncogene Wnt-3 Antibody, Int 4 Antibody, WNT3 Antibody, wnt3 Antibody

Immunogen

Synthetic peptide from the C-terminal of Human Proto-oncogene Wnt-3 (aa. 280-330)

Purification

Peptide Affinity Purified

Storage **-20°C**

Storage Buffer

PBS pH 7.4, 50% glycerol, 0.09% sodium azide *Storage buffer may change when conjugated

Shipping Temperature

Blue Ice or 4°C

Certificate of Analysis

A 1:1000 dilution of SPC-783 was sufficient for detection of WNT3 in 15 µg of mouse Brain cell lysates by ECL immunoblot analysis using goat anti-rabbit IgG:HRP as the secondary antibody.

Cellular Localization


Secreted | Extracellular Space | Extracellular Matrix

WNT3 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

WNT3 Antibody - Images


Immunohistochemistry analysis using Rabbit Anti-WNT3 Polyclonal Antibody (SPC-783). Tissue: Breast. Species: Human. Fixation: Formalin Fixed Paraffin-Embedded. Primary Antibody: Rabbit Anti-WNT3 Polyclonal Antibody (SPC-783) at 1:50 for 30 min at RT. Counterstain: Hematoxylin. Magnification: 10X. HRP-DAB Detection. | Western blot analysis of Mouse Brain showing detection of 39.6 kDa WNT3 protein using Rabbit Anti-WNT3 Polyclonal Antibody (SPC-783). Load: 15 μ g. Block: 5% Skim Milk powder in TBST. Primary Antibody: Rabbit Anti-WNT3 Polyclonal Antibody (SPC-783) at 1:1000 for 2 hours at RT with shaking. Secondary Antibody: Goat Anti-Rabbit IgG: HRP at 1:5000 for 1 hour at RT. Color Development: ECL solution for 5 min at RT. Predicted/Observed Size: 39.6 kDa. Other Band(s): 50 kDa.



Immunohistochemistry analysis using Rabbit Anti-WNT3 Polyclonal Antibody (SPC-783). Tissue: Breast. Species: Human. Fixation: Formalin Fixed Paraffin-Embedded. Primary Antibody: Rabbit Anti-WNT3 Polyclonal Antibody (SPC-783) at 1:50 for 30 min at RT. Counterstain: Hematoxylin. Magnification: 10X. HRP-DAB Detection. | Western blot analysis of Mouse Brain showing detection of 39.6 kDa WNT3 protein using Rabbit Anti-WNT3 Polyclonal Antibody (SPC-783). Load: 15 μ g. Block: 5% Skim Milk powder in TBST. Primary Antibody: Rabbit Anti-WNT3 Polyclonal Antibody (SPC-783) at 1:1000 for 2 hours at RT with shaking. Secondary Antibody: Goat Anti-Rabbit IgG: HRP at 1:5000 for 1 hour at RT. Color Development: ECL solution for 5 min at RT. Predicted/Observed Size: 39.6 kDa. Other Band(s): 50 kDa.