

**NRAS Antibody**  
**Rabbit Anti-Human NRAS Polyclonal**  
**Catalog # ASM10645**

**Specification**

---

**NRAS Antibody - Product Information**

|                           |                             |
|---------------------------|-----------------------------|
| Application               | <b>WB</b>                   |
| Primary Accession         | <a href="#">P01111</a>      |
| Other Accession           | <a href="#">NP_002515.1</a> |
| Host                      | <b>Rabbit</b>               |
| Reactivity                | <b>Rat</b>                  |
| Clonality                 | <b>Polyclonal</b>           |
| Format                    | <b>NRAS</b>                 |
| <b>Target/Specificity</b> |                             |
| NRAS                      |                             |

**Other Names**

Transforming protein N-Ras Antibody, GTPase NRas Antibody, NRAS Antibody, HRAS1 Antibody

**Immunogen**

Synthetic peptide from the C-terminal of Human GTPase NRas (aa. 85-185)

**Purification**

Peptide Affinity Purified

Storage **-20°C**

**Storage Buffer**

PBS pH 7.4, 50% glycerol, 0.09% sodium azide \*Storage buffer may change when conjugated

Shipping Temperature

**Blue Ice or 4°C**

**Certificate of Analysis**

A 1:1000 dilution of SPC-796 was sufficient for detection of NRAS in 15 µg of rat Kidney cell lysates by ECL immunoblot analysis using goat anti-rabbit IgG:HRP as the secondary antibody.


**Cellular Localization**

Cell Membrane | Lipid-Anchor | Cytoplasmic Side | Golgi Apparatus Membrane | Lipid-Anchor

**NRAS Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**NRAS Antibody - Images**

Western blot analysis of Rat Kidney showing detection of 21.2 kDa NRAS protein using Rabbit Anti-NRAS Polyclonal Antibody (SPC-796). Load: 15  $\mu$ g. Block: 5% Skim Milk powder in TBST. Primary Antibody: Rabbit Anti-NRAS Polyclonal Antibody (SPC-796) at 1:1000 for 2 hours at RT with shaking. Secondary Antibody: Goat Anti-Rabbit IgG: HRP at 1:5000 for 1 hour at RT. Color Development: ECL solution for 5 min at RT. Predicted/Observed Size: 21.2 kDa. Other Band(s): 25 kDa.