

**VPS35 Antibody**  
**VPS35 Antibody, Clone 8A3**  
**Catalog # ASM10693**

**Specification**

---

**VPS35 Antibody - Product Information**

Application	<b>IHC, IP, WB</b>
Primary Accession	<a href="#">O96OK1</a>
Other Accession	<a href="#">NP_060676.2</a>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
<b>Target/Specificity</b>	
VPS35	

**Other Names**

Vacuolar protein sorting-associated protein 35, MEM3, PARK17, VPS35 retromer complex component, maternal-embryonic 3, vesicle protein sortin 35, TCCCTA00141, FLJ10752

**Immunogen**

Full length recombinant human VSP35

**Purification**

Protein G Purified

Storage **-20°C**

**Storage Buffer**

PBS pH 7.4, 50% glycerol, 0.09% Sodium azide \*Storage buffer may change when conjugated

Shipping Temperature **Blue Ice or 4°C**

**Certificate of Analysis**

A 1:1000 dilution of SMC-604 was sufficient for detection of VPS35 in 10 µg of SH-SY5Y by ECL immunoblot analysis using Goat Anti-Mouse IgG:HRP as the secondary antibody.

**Cellular Localization**

Endosome | Lysosome | Vesicles | Cytoplasm | Membrane


**VPS35 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**VPS35 Antibody - Images**







---




Immunohistochemistry analysis using Mouse Anti-VPS35 Monoclonal Antibody, Clone 8A3 (ASM10693). Tissue: Kidney. Species: Mouse. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10693) at 1:100 for Overnight at 4C, then 30 min at 37C. Secondary Antibody: Goat Anti-Mouse IgG (H+L): FITC for 45 min at 37C. Counterstain: DAPI for 3 min at RT. Magnification: 20X.




Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-VPS35 Monoclonal Antibody, Clone 8A3 (ASM10693). Tissue: A549 cells. Species: Human. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10693) at 1:5 (tissue culture supernatant). Secondary Antibody: Donkey anti-mouse: Alexa Fluor 594 at 1:4000 in 0.2% BSA PBS. Counterstain: DAPI. Localization: Vesicles. A) VPS35 KO A549 cells B) WT A549 cells. Courtesy of: Dario Alessi Lab, University of Dundee.

VPS35 Antibody Concentration	SMC-604 (8A3)	
	A549 WT	A549 VPS35 KO
1/500		
1/1000		
1/2000		


Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-VPS35 Monoclonal Antibody, Clone 8A3 (ASM10693). Tissue: A549 WT, VPS35 KO cells. Species: Human. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10693). Secondary Antibody: Donkey Anti-Mouse AlexaFluor 594. Clone can detect VPS35 at 1/2000 concentration.




Immunoprecipitation analysis using Mouse Anti-VPS35 Monoclonal Antibody, Clone 8A3 (ASM10693). Tissue: A549 cells. Species: Human. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10693). 500  $\mu$ L cell culture supernatants were incubated with 10  $\mu$ L of Protein A/G resin beads for 1 hour at 4°C. ASM10693 clone 8A3 depletes VPS35 from the A549 cell extract..




Immunoprecipitation analysis using Mouse Anti-VPS35 Monoclonal Antibody, Clone 8A3 (ASM10693). Tissue: A549 cells. Species: Human. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10693).




Western Blot analysis of Human, Mouse A549, MEF showing detection of VPS35 protein using Mouse Anti-VPS35 Monoclonal Antibody, Clone 8A3 (ASM10693). Lane 1: Molecular Weight Ladder. Lane 2: VPS35 KO A549 cells. Lane 3: mouse embryonic fibroblast cells.. Load: 8  $\mu$ g each A549 and MEF. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10693) at 1:5 (tissue culture supernatant). Secondary Antibody: Donkey anti-mouse IRDye 800CW at 1:25000 in TBS-T.



Western Blot analysis of Human SH-SY5Y showing detection of VPS35 protein using Mouse Anti-VPS35 Monoclonal Antibody, Clone 8A3 (ASM10693). Lane 1: Molecular Weight Ladder. Lane 2: SH-SY5Y (10  $\mu$ g). Load: 10  $\mu$ g. Block: 5% Skim Milk powder in TBST. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10693) at 1:1000 for 2 hours at RT with shaking. Secondary Antibody: Goat anti-mouse IgG:HRP at 1:4000 for 1 hour at RT with shaking. Color Development: Chemiluminescent for HRP (Moss) for 5 min in RT.



Immunoprecipitation analysis using Mouse Anti-VPS35 Monoclonal Antibody, Clone 8A3 (ASM10693). Tissue: A549 cells. Species: Human. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10693). Three amounts of ASM10693 (3, 1 and 0.3 ug) were non-covalently coupled to 10uL of A/G sepharose beads for 1 hour at 4°C and next incubated with 250ug of A549 lysate for 2 hours at 4°C.



Immunoprecipitation analysis using Mouse Anti-VPS35 Monoclonal Antibody, Clone 8A3 (ASM10693). Tissue: embryonic fibroblast. Species: Mouse. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10693). Three amounts of ASM10693 (3, 1 and 0.3 ug) were non-covalently coupled to 10uL of A/G sepharose beads for 1 hour at 4°C and next incubated with 250ug of MEF lysate for 2 hours at 4°C.

### VPS35 Antibody - Background

Vacuolar Protein Sorter-35 (VPS35) is a component of the retromer complex, which is essential for endosome-to-Golgi retrieval of membrane proteins. VPS35 mutations such as D620N have been linked to Parkinson's Disease (PD) (1,2) and affect retromer function, protein homeostasis, and mitochondria (3).

### VPS35 Antibody - References

1. Vilarino-Guell, C. et al. (2011) Am J Hum Genet 89:162-167

2. Zimprich, A. et al. (2011) Am J Hum Genet 89:168-175
3. Rahman, A.A., Morrison, B.E. (2019) Neurosci 401:1-10.