

# Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody Goat Polyclonal, Fluorescein (FITC)

Catalog # ASR1615

### **Specification**

**Target Species** 

**Physical State** 

Target Isotype

Host Isotype

Immunogen

Reconstitution Volume

Reconstitution Buffer

Buffer

Clonality

## Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody - Product Information

Description Anti-MOUSE IgG (H&L) (GOAT) Antibody

Fluorescein Conjugated (Min X Human

**Serum Proteins**)

Host Goat

Conjugate Fluorescein (FITC)

FP Value 6.6 moles Fluorescein (FITC) per mole of

IgG Mouse Polyclonal

Application ,3,4,
Application Note IF Microscopy

1:500-1:2,500;FlowCytometry

1:1,000-1:5,000 Lyophilized

IgG

IgG (H&L)

0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2
Mouse IgG whole molecule

1.0 mL

Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

## Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody - Additional Information

### **Shipping Condition**

**Ambient** 

### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Goat Serum, Mouse IgG and Mouse Serum. No reaction was observed against Human Serum Proteins.

#### **Storage Condition**

Store secondary antibody at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at



4° C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information

Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody - Images

Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody - Background

Mouse secondary antibody conjugated to FITC is available in a variety of formats. Anti IgG secondary antibody conjugated is suitable for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. Anti mouse IgG FITC is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody -

Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody Citations

• Therapeutic effects of acylated ghrelin-specific receptor GHS-R1a antagonist in islet transplantation