

**Anti-Mouse Red Blood Cell RBC Secondary Antibody**  
**Rabbit Polyclonal, Unconjugated**  
**Catalog # ASR2645**

**Specification**

---

**Anti-Mouse Red Blood Cell RBC Secondary Antibody - Product Information**

Description	<b>Anti-MOUSE Red Blood Cell (RBC) (RABBIT) Antibody</b>
Host	<b>Rabbit</b>
Conjugate	<b>Unconjugated</b>
Clonality	<b>Polyclonal</b>
Application	<b>''</b>
Application Note	<b>AGGLUTINATION TITER 1:32-1:64</b>
Physical State	<b>Lyophilized</b>
Host Isotype	<b>Antiserum</b>
Buffer	<b>0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>
Immunogen	<b>Mouse washed pooled Red Blood Cells (RBC)</b>
Species of Origin	<b>Mouse</b>
Reconstitution Volume	<b>2.0 mL</b>
Reconstitution Buffer	<b>Restore with deionized water (or equivalent)</b>
Stabilizer	<b>None</b>
Preservative	<b>0.01% (w/v) Sodium Azide</b>

**Anti-Mouse Red Blood Cell RBC Secondary Antibody - Additional Information**

**Shipping Condition**

Ambient

**Purity**

This product was prepared from polyspecific antiserum by a delipidation and defibrination.

**Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**Anti-Mouse Red Blood Cell RBC Secondary Antibody - Protein Information**

**Anti-Mouse Red Blood Cell RBC Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Anti-Mouse Red Blood Cell RBC Secondary Antibody - Images**

#### **Anti-Mouse Red Blood Cell RBC Secondary Antibody - Citations**

- [Hydrogen sulfide donor NaHS alters antibody structure and function via sulfhydration.](#)