

Anti-Cyclin E (RABBIT) Antibody Cyclin E Antibody Catalog # ASR3664

Specification

Anti-Cyclin E (RABBIT) Antibody - Product Information

an clonal E, IP, I, LCI -Cyclin-E antibody is suitable for the ction by immunoblot, ELISA and IP. earchers should determine optimal s for applications that are not stated w. Antiserum will specifically react a 45 kDa Cyclin E protein from human
ie. id (sterile filtered) -Cyclin-E Antibody was produced by ated immunizations with a KLH ugated peptide corresponding to a on near the carboxy terminus of human in E. % (w/v) Sodium Azide

Anti-Cyclin E (RABBIT) Antibody - Additional Information

Gene ID 898

Other Names 898

Purity

Anti-Cyclin-E was prepared from monospecific antiserum by delipidation and defibrination. No reaction was observed against other related cyclins. Cross reactivity with Cyclin E from other species may also occur.

Storage Condition

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Cyclin E (RABBIT) Antibody - Protein Information



Name CCNE1

Synonyms CCNE

Function Essential for the control of the cell cycle at the G1/S (start) transition.

Cellular Location Nucleus.

Tissue Location Highly expressed in testis and placenta. Low levels in bronchial epithelial cells.

Anti-Cyclin E (RABBIT) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Cyclin E (RABBIT) Antibody - Images

Anti-Cyclin E (RABBIT) Antibody - Background

Anti-Cyclin-E antibody detects cyclin-E. Cyclin E is a member of the cyclin family. Cyclin E binds to the G1 phase Cdk2, which is required for the transition from G1 to S phase of the cell cycle that determines cell division. The Cyclin E/CDK2 complex phosphorylates p27Kip1, tagging it for degradation, thus promoting expression of Cyclin A, allowing progression to S phase. Anti-cyclin-e Antibody is ideal for investigators involved in Cell Signaling, cell biology and Signal Transduction research.