

**Anti-SKP1 (C-terminal specific) (RABBIT) Antibody**  
**SKP1 Antibody**  
**Catalog # ASR3721****Specification**

---

**Anti-SKP1 (C-terminal specific) (RABBIT) Antibody - Product Information**

Host	Rabbit
Conjugate	Unconjugated
Target Species	Human
Reactivity	Human
Clonality	Polyclonal
Application	WB, IHC, E, I, LCI
Application Note	This antibody reacts with human SKP1 by western blot and immunoprecipitation. The antibody immunoprecipitates in vitro translated protein and protein from cell lysates (using HeLa, NIH-3T3, and others). Coimmunoprecipitation of cyclin A and Cuo1 may occur. An 18.5 kDa band corresponding to human SKP1 is detected. Most cell lines expressing SKP1 can be used as a positive control. Researchers should determine optimal titers for other applications.
Physical State	Liquid (sterile filtered)
Immunogen	This antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to amino acids 152-163 of Human SKP1 (C-terminus) coupled to KLH.
Preservative	0.01% (w/v) Sodium Azide

**Anti-SKP1 (C-terminal specific) (RABBIT) Antibody - Additional Information****Gene ID** 6500**Other Names**  
6500**Purity**

This product is monospecific antiserum processed by delipidation and defibrination followed by sterile filtration. This product reacts with human SKP1. Cross reactivity is expected against mouse SKP1 based on a high degree of sequence homology. Cross reactivity with other SKP1 proteins or SKP1 from other sources is not known.

**Storage Condition**

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

### Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

## Anti-SKP1 (C-terminal specific) (RABBIT) Antibody - Protein Information

**Name** SKP1

**Synonyms** EMC19, OCP2, SKP1A, TCEB1L

### Function


Essential component of the SCF (SKP1-CUL1-F-box protein) ubiquitin ligase complex, which mediates the ubiquitination of proteins involved in cell cycle progression, signal transduction and transcription. In the SCF complex, serves as an adapter that links the F-box protein to CUL1. The functional specificity of the SCF complex depends on the F-box protein as substrate recognition component. SCF(BTRC) and SCF(FBXW11) direct ubiquitination of CTNNB1 and participate in Wnt signaling. SCF(FBXW11) directs ubiquitination of phosphorylated NFKBIA. SCF(BTRC) directs ubiquitination of NFKBIB, NFKBIE, ATF4, SMAD3, SMAD4, CDC25A, FBXO5, CEP68 and probably NFKB2 (PubMed:<a href="http://www.uniprot.org/citations/25704143" target="\_blank">25704143</a>). SCF(SKP2) directs ubiquitination of phosphorylated CDKN1B/p27kip and is involved in regulation of G1/S transition. SCF(SKP2) directs ubiquitination of ORC1, CDT1, RBL2, ELF4, CDKN1A, RAG2, FOXO1A, and probably MYC and TAL1. SCF(FBXW7) directs ubiquitination of cyclin E, NOTCH1 released notch intracellular domain (NICD), and probably PSEN1. SCF(FBXW2) directs ubiquitination of GCM1. SCF(FBXO32) directs ubiquitination of MYOD1. SCF(FBXO7) directs ubiquitination of BIRC2 and DLGAP5. SCF(FBXO33) directs ubiquitination of YBX1. SCF(FBXO11) directs ubiquitination of BCL6 and DTL but does not seem to direct ubiquitination of TP53. SCF(BTRC) mediates the ubiquitination of NFKBIA at 'Lys-21' and 'Lys-22'; the degradation frees the associated NFKB1-RELA dimer to translocate into the nucleus and to activate transcription. SCF(CCNF) directs ubiquitination of CCP110. SCF(FBXL3) and SCF(FBXL21) direct ubiquitination of CRY1 and CRY2. SCF(FBXO9) directs ubiquitination of TTI1 and TELO2. SCF(FBXO10) directs ubiquitination of BCL2. Core component of the Cul7-RING(FBXW8) ubiquitin ligase complex, which mediates the ubiquitination and subsequent proteasomal degradation of target proteins (PubMed:<a href="http://www.uniprot.org/citations/35982156" target="\_blank">35982156</a>).


## Anti-SKP1 (C-terminal specific) (RABBIT) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-SKP1 (C-terminal specific) (RABBIT) Antibody - Images





Most modifiers mature by proteolytic processing from inactive precursors (a; amino acid). Arrowheads point to the cleavage sites. Ubiquitin is expressed either as polyubiquitin or as a fusion with ribosomal proteins. Conjugation requires activating (E1) and conjugating (E2) enzymes that form thioesters (S) with the modifiers. Modification of cullins by RUB involves SCF(SKP1/cullin-1/F-box protein) /CBC(cullin-2/elongin B/elonginC) -like E3 enzymes that are also involved in ubiquitination. In contrast to ubiquitin, the UBLs do not seem to form multi-UBL chains. UCRP(ISG15) resembles two ubiquitin moieties linked head-to-tail. Whether HUB1 functions as a modifier is currently unclear. APG12 and URM1 are distinct from the other modifiers because they are unrelated in sequence to ubiquitin. Data contributed by S.Jentsch.

### Anti-SKP1 (C-terminal specific) (RABBIT) Antibody - Background

SKP1 is also known as S-phase kinase-associated protein 1A, Cyclin A/CDK2-associated protein p19, p19A, p19skp1, RNA polymerase II elongation factor-like protein, Organ of Corti protein 2, OCP-II protein, OCP-2, Transcription elongation factor B, and SIII. SKP1 is an essential component of the SCF (SKP1-CUL1-F-box protein) ubiquitin ligase complex, which mediates the ubiquitination of proteins involved in cell cycle progression, signal transduction and transcription. In the SCF complex, SKP1 serves as an adapter that links the F-box protein to CUL1. SKP1 interacts directly with CUL1 and F-box proteins, such as BTRC and SKP2, in the SCF complex. SKP1 also interacts with FBX29 and the cyclin A/CDK2 complex. SKP1 is part of a SCF-like complex consisting of CUL7, RBX1, SKP1 and FBXW8 and is also a component of a E3 ubiquitin ligase complex containing UBE2D1, SIAH1, CACYBP/SIP, SKP1A, APC and TBL1X.