

Anti-IK α C-terminal (RABBIT) Antibody
IKB alpha Antibody
Catalog # ASR3805

Specification

Anti-IK α C-terminal (RABBIT) Antibody - Product Information

Host	Rabbit
Conjugate	Unconjugated
Target Species	Human
Reactivity	Rat, Human, Mouse
Clonality	Polyclonal
Application	WB, IHC, E, I, LCI
Application Note	Anti-IkBa Antibody has been tested by western blot and is suitable for ELISA, and IHC. Researchers should determine optimal titers for applications that are not stated below.
Physical State	Liquid (sterile filtered)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	IKB alpha Antibody was produced by repeated immunizations with a synthetic IkBa peptide corresponding to a region near the C-terminus of the human protein.
Preservative	0.1% (w/v) Sodium Azide

Anti-IK α C-terminal (RABBIT) Antibody - Additional Information

Gene ID 4792

Other Names
4792

Purity

IKB alpha Antibody was prepared from monospecific antiserum by delipidation and defibrination. Anti- IkBa may react non-specifically with other proteins. Control peptide (code #100-4167P) will compete only with the specific reaction of antiserum with IkBa.

Storage Condition

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-IK α C-terminal (RABBIT) Antibody - Protein Information

Name NFKBIA

Synonyms IKBA, MAD3, NFKBI

Function

Inhibits the activity of dimeric NF-kappa-B/REL complexes by trapping REL (RELA/p65 and NFKB1/p50) dimers in the cytoplasm by masking their nuclear localization signals (PubMed:1493333, PubMed:36651806, PubMed:7479976). On cellular stimulation by immune and pro-inflammatory responses, becomes phosphorylated promoting ubiquitination and degradation, enabling the dimeric RELA to translocate to the nucleus and activate transcription (PubMed:7479976, PubMed:7628694, PubMed:7796813, PubMed:7878466).

Cellular Location


Cytoplasm. Nucleus. Note=Shuttles between the nucleus and the cytoplasm by a nuclear localization signal (NLS) and a CRM1-dependent nuclear export.

Anti-IKBA C-terminal (RABBIT) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IKBA C-terminal (RABBIT) Antibody - Images



Western Blot of Rabbit anti-IKBA C-terminal antibody. Lane 1: IKBA C-terminal. Load: 50 ng per lane. Primary antibody: IKBA C-terminal antibody at 1:500 for overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody at 1:40,000 for 30 min at RT. Block: Blocking Buffer for Fluorescent Western Blotting (MB-070) for 30 min RT. Predicted/Observed size: 61 kDa,

61 kDa for IKB α C-terminal.

Anti-IKB α C-terminal (RABBIT) Antibody - Background

Anti-IKB alpha Antibody detects Ikb α . I-kappa-B-alpha inhibits the activity of dimeric NF-kappa-B/REL complexes by trapping REL dimers in the cytoplasm through masking of their nuclear localization signals. On cellular stimulation by immune and proinflammatory responses, IKBA becomes phosphorylated promoting ubiquitination and degradation, enabling the dimeric RELA to translocate to the nucleus and activate transcription. Anti-IKB α Antibody is ideal for investigators involved in Cell Signaling, Neuroscience, Signal Transduction and Immunology research.