

Anti-ACETYL CHOLINESTERASE (Bovine Erythrocytes) (GOAT) Antibody
Acetyl Cholinesterase Antibody
Catalog # ASR3938**Specification****Anti-ACETYL CHOLINESTERASE (Bovine Erythrocytes) (GOAT) Antibody - Product Information**

Host	Goat
Conjugate	Unconjugated
Target Species	Bovine
Reactivity	Bovine
Clonality	Polyclonal
Application	WB, E, IP, I, LCI
Application Note	Anti-Acetyl Cholinesterase Antibody has been tested by western blot and is suitable to be assayed against 1.0 µg of Acetyl Cholinesterase [Bovine Erythrocytes] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] (Rabbit) code #605-4302 and (ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:1,200 to 1:5,000 of the reconstitution concentration is suggested for this product.
Physical State	Lyophilized
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Acetyl Cholinesterase [Bovine Erythrocytes]
Reconstitution Volume	100 µL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Preservative	0.01% (w/v) Sodium Azide

Anti-ACETYL CHOLINESTERASE (Bovine Erythrocytes) (GOAT) Antibody - Additional Information**Gene ID** 540446**Other Names**
540446**Purity**

Anti-Acetyl Cholinesterase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum as well as purified and partially purified Acetyl Cholinesterase [Bovine Erythrocytes]. Cross reactivity against

Acetyl Cholinesterase from other sources is unknown.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-ACETYL CHOLINESTERASE (Bovine Erythrocytes) (GOAT) Antibody - Protein Information

Name ACHE

Function

Terminates signal transduction at the neuromuscular junction by rapid hydrolysis of the acetylcholine released into the synaptic cleft.

Cellular Location

Synapse. Secreted. Cell membrane; Peripheral membrane protein

Anti-ACETYL CHOLINESTERASE (Bovine Erythrocytes) (GOAT) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-ACETYL CHOLINESTERASE (Bovine Erythrocytes) (GOAT) Antibody - Images**Anti-ACETYL CHOLINESTERASE (Bovine Erythrocytes) (GOAT) Antibody - Background**

Acetylcholinesterase terminates signal transduction at the neuromuscular junction by rapid hydrolysis of the acetylcholine released into the synaptic cleft.