

Anti-LACTATE DEHYDROGENASE (Rabbit Muscle) (GOAT) Antibody Peroxidase Conjugated**Lactate Dehydrogenase Antibody Peroxidase Conjugated
Catalog # ASR4098****Specification**

Anti-LACTATE DEHYDROGENASE (Rabbit Muscle) (GOAT) Antibody Peroxidase Conjugated - Product Information

Host	Goat
Conjugate	Peroxidase (Horseradish)
Target Species	Rabbit
Reactivity	Rabbit
Clonality	Polyclonal
Application	WB, IHC, E, IP, I, LCI
Application Note	Anti-Lactate Dehydrogenase peroxidase Antibody has been tested in ELISA, western blot, and immunofluorescence. This product is suitable for IP and IHC. Researchers should determine optimal titers for applications that are not stated below.
Physical State	Lyophilized
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Anti-Lactate Dehydrogenase Antibody was produced by repeated immunizations with rabbit muscle Lactate Dehydrogenase full length protein.
Reconstitution Volume	100 µL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Anti-LACTATE DEHYDROGENASE (Rabbit Muscle) (GOAT) Antibody Peroxidase Conjugated - Additional Information**Gene ID** 100009107**Other Names**
100009107**Purity**

Anti-Lactate Dehydrogenase Antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum as well as purified and partially purified Lactate Dehydrogenase [Rabbit Muscle]. Cross reactivity

against Lactate Dehydrogenase from other tissues and species may occur but have not been specifically determined.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-LACTATE DEHYDROGENASE (Rabbit Muscle) (GOAT) Antibody Peroxidase Conjugated - Protein Information

Name LDHA

Function

Interconverts simultaneously and stereospecifically pyruvate and lactate with concomitant interconversion of NADH and NAD(+).

Cellular Location

Cytoplasm.

Anti-LACTATE DEHYDROGENASE (Rabbit Muscle) (GOAT) Antibody Peroxidase Conjugated - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-LACTATE DEHYDROGENASE (Rabbit Muscle) (GOAT) Antibody Peroxidase Conjugated - Images

Anti-LACTATE DEHYDROGENASE (Rabbit Muscle) (GOAT) Antibody Peroxidase Conjugated - Background

Anti-Lactate Dehydrogenase Antibody detects Lactate Dehydrogenase. Lactate dehydrogenases exist in four distinct enzyme classes. Two of them are cytochrome c-dependent enzymes, each acting on either D-lactate or L-lactate. The other two are NAD(P)-dependent enzymes, each acting on either D-lactate or L-lactate. Lactate dehydrogenase catalyzes the interconversion of pyruvate and lactate. Anti-Lactate Dehydrogenase Antibody is ideal for investigators involved in Cell Signaling, Immunology, Neuroscience and Signal Transduction research.