

Anti-GSTS1 (MOUSE) Monoclonal Antibody

GSTS1 Antibody Catalog # ASR4229

Specification

Physical State

Anti-GSTS1 (MOUSE) Monoclonal Antibody - Product Information

Host Mouse

Conjugate Unconjugated

Target Species Mouse
Reactivity Human
Clonality Monoclonal

Application WB, IHC, E, IP, I, LCI

Application Note Anti-GSTS1 antibody has been tested by

ELISA and Western Blot. Suitable for most immunological techniques requiring high titer binding and lot-to-lot consistency. Specific conditions for reactivity should be

optimized by the end user. Liquid (sterile filtered)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Immunogen This Protein A purified antibody was

prepared by repeated immunizations in

mice with S1 Protein

Preservative 0.01% (w/v) Sodium Azide

Anti-GSTS1 (MOUSE) Monoclonal Antibody - Additional Information

Purity

This product is purified from roller bottle culture by Protein A chromatography followed by extensive dialysis against the buffer stated above. Reacts specifically with S1 protein. Cross reactivity from other sources has not been determined.

Storage Condition

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-GSTS1 (MOUSE) Monoclonal Antibody - Protein Information


Anti-GSTS1 (MOUSE) Monoclonal Antibody - Protocols



Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-GSTS1 (MOUSE) Monoclonal Antibody - Images



Western Blot of Mouse anti-GSTS1 Monoclonal Antibody. Lane 1: rGSTS1 protein. Lane 2: GST (p/n 000-001-200). Load: 50 ng per lane. Primary antibody: Mouse anti-GSTS1 Monoclonal Antibody at 1:1,000 overnight at 4°C. Secondary antibody: Peroxidase conjugated Rb-a-Ms IgG (p/n 610-403-C46) at 1:40,000 for 30 min at RT. Block: 0.75% Casein-TBS (p/n MB-082) for 30 min at RT. Predicted/Observed size: 27 kDa, 27 kDa for GSTS1.

Anti-GSTS1 (MOUSE) Monoclonal Antibody - Background

Rockland produces a wide range of human GST antibodies in our laboratories. Select appropriate GST antibodies for your research by isotype, epitope, applications and species reactivity. There are 22 members of the human GST family of proteins. GST is responsible for the conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. The amino acid sequence GST is highly conserved in most organisms including mammals. GSTs proteins are typically homodimeric, with both heterologous GST dimers have been observed. GST monomers have an average molecular weight of approximately 25-28 kDa in size. Note a different form of non-human GST (Glutathione-S-Transferase) is used as a protein expression tag commonly in molecular biology applications. All anti-GST antibodies my not react with recombinant GST-fusion proteins.