

**Anti-6X HIS EPITOPE TAG (MOUSE) Monoclonal Antibody Peroxidase Conjugated**  
**6X His Tag Antibody HRP Conjugated**  
**Catalog # ASR4304****Specification**

---

**Anti-6X HIS EPITOPE TAG (MOUSE) Monoclonal Antibody Peroxidase Conjugated - Product Information**

Host	<b>Mouse</b>
Conjugate	<b>Peroxidase (Horseradish)</b>
Clonality	<b>Monoclonal</b>
Application	<b>WB, IHC, E, I, LCI</b>
Application Note	<b>Anti-6X His is optimally suited for monitoring expression of His-tagged fusion proteins. As such, anti-6X His/6X His can be used to identify fusion proteins containing the 6X His epitope. The antibody recognizes the His tag fused either to the amino- or carboxy- termini of targeted proteins. This antibody has been tested by ELISA and western blotting against both the immunizing peptide and His-containing recombinant proteins. Although not tested, this antibody is likely functional for immunoprecipitation and immunocytochemistry.</b>
Physical State	<b>Lyophilized</b>
Buffer	<b>0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>
Immunogen	<b>This 6X his antibody was produced in mice by repeated immunizations with 6X His epitope tag peptide H-H-H-H-H-H conjugated to KLH using maleimide.</b>
Reconstitution Volume	<b>100 µL</b>
Reconstitution Buffer	<b>Restore with deionized water (or equivalent)</b>
Stabilizer	<b>10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free</b>
Preservative	<b>0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!</b>

**Anti-6X HIS EPITOPE TAG (MOUSE) Monoclonal Antibody Peroxidase Conjugated - Additional Information****Purity**

6X His antibody is directed against the 6X His motif and was purified from concentrated tissue culture supernate by Protein A chromatography. This monoclonal anti-6X His tag antibody detects over-expressed proteins containing the 6X His epitope tag and is useful in determining its presence in various assays. To date, this antibody has reacted with all His tagged proteins so far tested. In western blotting of bacterial extracts, the antibody does not cross-react with endogenous proteins. The antibody recognizes the His-tag (His-His-His-His-His-His) fused to either

the amino- or carboxy-termini of targeted proteins in transfected or transformed cells.

### Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

### Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.


## Anti-6X HIS EPITOPE TAG (MOUSE) Monoclonal Antibody Peroxidase Conjugated - Protein Information

## Anti-6X HIS EPITOPE TAG (MOUSE) Monoclonal Antibody Peroxidase Conjugated - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-6X HIS EPITOPE TAG (MOUSE) Monoclonal Antibody Peroxidase Conjugated - Images



ELISA of Mouse anti-6xHIS Tag Antibody. Antigen: 6X HIS-tagged conjugated BSA at the N or C terminus of the 6XHis. Coating amount: 0.15ug per well. Primary antibody (direct detection): HRP conjugated 6xHIS Tag antibody diluted from stock concentration at 100ug/mL. Substrate: TMB (p/n TMBE-1000).

## Anti-6X HIS EPITOPE TAG (MOUSE) Monoclonal Antibody Peroxidase Conjugated - Background

Epitope tags are short peptide sequences that are easily recognized by tag-specific antibodies. Due to their small size, epitope tags do not affect the tagged protein's biochemical properties.

Most often, sequences encoding the epitope tag are included with target DNA at the time of cloning to produce fusion proteins containing the epitope tag sequence. This allows anti-epitope tag antibodies to serve as universal detection reagents for any tag-containing protein produced by recombinant means. This means that anti-epitope tag antibodies are a useful alternative to generating specific antibodies to identify, immunoprecipitate or immunoaffinity purify a recombinant protein. The anti-epitope tag antibody is usually functional in a variety of antibody-dependent experimental procedures. Expression vectors producing epitope tag fusion proteins are available for a variety of host expression systems including bacteria, yeast, insect and mammalian cells. Rockland Immunochemicals produces anti-epitope tag antibodies against many common epitope tags including Myc, GST, GFP, 6X His, MBP, FLAG and HA. Rockland Immunochemicals also produces antibodies to other tags including FITC, Rhodamine (TRITC), DNP and biotin.