

Anti-6X HIS EPITOPE TAG (MOUSE) Monoclonal ATTO 647N Conjugated Antibody

6X His Tag ATTO 647N Conjugated Antibody Catalog # ASR4341

Specification

Anti-6X HIS EPITOPE TAG (MOUSE) Monoclonal ATTO 647N Conjugated Antibody - Product Information

Host Mouse
Conjugate ATTO 647N
FP Value 1.0
Clonality Monoclonal

Application WB, I, LCI

Application Note

Anti-6X His is optimally suited for monitoring expression of His-tagged fusion proteins. As such, anti-6X His/6X His can

be used to identify fusion proteins that contain the 6X His epitope. The antibody recognizes the His tag fused either to the amino- or carboxy- termini of targeted proteins. This antibody has been tested by ELISA, dot blot, and western blotting against both the immunizing peptide and His-containing recombinant proteins. ATTOs are designed for STED microscopy, FRET, immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex

analysis, including multicolor imaging, utilizing various commercial platforms.

Physical State

Lyophilized

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Immunogen HIS Tag antibody was produced in mice by

repeated immunizations with 6X His epitope tag peptide H-H-H-H-H conjugated to KLH using maleimide.

Reconstitution Volume 50 µL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -


Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Anti-6X HIS EPITOPE TAG (MOUSE) Monoclonal ATTO 647N Conjugated Antibody - Additional Information

Purity

6X HIS Epitope Tag ATTO 647N conjugated antibody is directed against the 6X His motif and is useful in determining its presence in various assays. This monoclonal anti-6X His tag antibody





detects over-expressed proteins containing the 6X His epitope tag. To date, this antibody has reacted with all His tagged proteins so far tested. In western blotting of bacterial extracts, the antibody does not cross-react with endogenous proteins. The antibody recognizes the His-tag (His-His-His-His-His-His) fused to either the amino- or carboxy-termini of targeted proteins in transfected or transformed cells.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-6X HIS EPITOPE TAG (MOUSE) Monoclonal ATTO 647N Conjugated Antibody - Protein Information

Anti-6X HIS EPITOPE TAG (MOUSE) Monoclonal ATTO 647N Conjugated Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-6X HIS EPITOPE TAG (MOUSE) Monoclonal ATTO 647N Conjugated Antibody - Images

Anti-6X HIS EPITOPE TAG (MOUSE) Monoclonal ATTO 647N Conjugated Antibody - Background

6X His Tag ATTO 647N conjugated Antibody as well as other Epitope tags are short peptide sequences that are easily recognized by tag-specific antibodies. Due to their small size, epitope tags do not affect the tagged protein's biochemical properties. Most often sequences encoding the epitope tag are included with target DNA at the time of cloning to produce fusion proteins containing the epitope tag sequence. This allows anti-epitope tag antibodies to serve as universal detection reagents for any tag containing protein produced by recombinant means. This means that anti-epitope tag antibodies are a useful alternative to generating specific antibodies to identify, immunoprecipitate or immunoaffinity purify a recombinant protein. The anti-epitope tag antibody is usually functional in a variety of antibody-dependent experimental procedures. Expression vectors producing epitope tag fusion proteins are available for a variety of host expression systems including bacteria, yeast, insect and mammalian cells. Rockland Immunochemicals produces anti-epitope tag antibodies against many common epitope tags including Myc, GST, GFP, 6X His, MBP, FLAG and HA. Rockland Immunochemicals also produces antibodies to other tags including FITC, Rhodamine (TRITC), DNP and biotin.