

Anti-RFX5 (N-terminal) (RABBIT) Antibody

RFX5 Antibody Catalog # ASR4383

Specification

Anti-RFX5 (N-terminal) (RABBIT) Antibody - Product Information

Host Rabbit

Conjugated Unconjugated

Target Species
Reactivity
Human
Clonality
Application
Human
Polyclonal
WB, E, I, LCI

Application Note This product was assayed by immunoblot and found to be reactive against RFX5

(N-terminal specific) from a variety of fibroblast and B-cell lysates at a dilution of 1:500 followed by reaction with Peroxidase conjugated Affinity Purified anti-Rabbit IgG [H&L] (Goat) code #611-1302. Anti-RFX5 (N-terminal specific) is suitable for the detection by immunoblot of human RFX5 a shows a 75 kDa band. This product was also tested in a gel supershift assay and

found to be reactive against RFX5 complexes using 0.5 to 1.0 µl per assay.

Liquid (sterile filtered)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Immunogen RFX5 (N-terminal specific) peptide

corresponding to a region near the N-terminus of the human protein

conjugated to Keyhole Limpet Hemocyanin

(KLH).

Preservative 0.01% (w/v) Sodium Azide

Anti-RFX5 (N-terminal) (RABBIT) Antibody - Additional Information

Gene ID 5993

Physical State

Other Names

5993

Purity

This product was prepared from monospecific antiserum by delipidation, salt fractionation and ion exchange chromatography. Anti-RFX5 (N-terminal specific) may react with unknown minor bands at 55 kDa and 85 kDa.

Storage Condition

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after



Tel: 858.875.1900 Fax: 858.875.1999

standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-RFX5 (N-terminal) (RABBIT) Antibody - Protein Information

Name RFX5

Function

Activates transcription from class II MHC promoters. Recognizes X-boxes. Mediates cooperative binding between RFX and NF-Y. RFX binds the X1 box of MHC-II promoters.

Cellular Location

Nucleus.

Tissue Location

Ubiquitous.

Anti-RFX5 (N-terminal) (RABBIT) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-RFX5 (N-terminal) (RABBIT) Antibody - Images

Anti-RFX5 (N-terminal) (RABBIT) Antibody - Background

DNA-binding protein RFX5 activates transcription from class II MHC promoters and recognizes X-boxes. It mediates cooperative binding between RFX and NF-Y. RFX binds the X1 box of MHC-II promoters. RFX5 may be associated with Bare Lymphocyte Syndrome Type Ii Complementation group C and Mhc Class Ii Deficiency. Anti-RFX5 Antibody is useful for researchers interested in Transcription factors, Tuberculosis, and Allograft rejection.