

## Anti-Macrophage migration inhibitory factor (MIF) (CHICKEN) Antibody

MIF Antibody

Catalog # ASR4847

### Specification

---

#### Anti-Macrophage migration inhibitory factor (MIF) (CHICKEN) Antibody - Product Information

Host	Chicken
Conjugate	Unconjugated
Target Species	Human
Reactivity	Human
Clonality	Polyclonal
Application	WB, E, I, LCI
Application Note	This IgY fraction antibody of anti-Human macrophage migration inhibitory factor (MIF) has been tested for use in ELISA and immunoblotting. Although not tested, this antibody may also be useful for neutralization assays, immunohistochemistry and flow cytometry. The antibody recognizes 12,300 MW mature human MIF. The MIF gene encodes a protein of 115 aa. The initiating methionine is cleaved leaving a mature protein of 114 aa. Reactivity in other immunoassays is unknown.
Physical State	Lyophilized
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	This IgY fraction antibody was prepared from eggs of chickens laid after repeated immunizations with a synthetic peptide corresponding to aa 2-32 of Human MIF conjugated to keyhole limpet hemocyanin (KLH). MIF is a proinflammatory cytokine that plays an important role in systemic inflammatory events.
Reconstitution Volume	500 µL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Preservative	0.01% (w/v) Sodium Azide

#### Anti-Macrophage migration inhibitory factor (MIF) (CHICKEN) Antibody - Additional Information

Gene ID 4282

Other Names

4282

**Purity**

This product is an IgY fraction antibody purified from monospecific chicken egg yolks by a multi-step process which includes selective precipitation and salt fractionation followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Chicken Serum. The antibody is directed against the 12,300 MW human MIF protein and is useful in determining its presence in various assays. If neutralization experiments are performed for human MIF activity in bioassays, it is recommended to incubate the sample with a 1:500 dilution of the antibody for at least 4 hours before being tested. A control of similarly diluted normal chicken IgG is recommended. If FACS analysis experiments are performed for human MIF caution should be exhibited as the Fc domain of the chicken IgG molecule may interact with cells non-specifically.

**Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**Anti-Macrophage migration inhibitory factor (MIF) (CHICKEN) Antibody - Protein Information**

**Name** MIF {ECO:0000303|PubMed:2552447, ECO:0000312|HGNC:HGNC:7097}

**Function**

Pro-inflammatory cytokine involved in the innate immune response to bacterial pathogens (PubMed:<a href="http://www.uniprot.org/citations/15908412" target="\_blank">15908412</a>, PubMed:<a href="http://www.uniprot.org/citations/17443469" target="\_blank">17443469</a>, PubMed:<a href="http://www.uniprot.org/citations/23776208" target="\_blank">23776208</a>). The expression of MIF at sites of inflammation suggests a role as mediator in regulating the function of macrophages in host defense (PubMed:<a href="http://www.uniprot.org/citations/15908412" target="\_blank">15908412</a>, PubMed:<a href="http://www.uniprot.org/citations/17443469" target="\_blank">17443469</a>, PubMed:<a href="http://www.uniprot.org/citations/23776208" target="\_blank">23776208</a>). Counteracts the anti-inflammatory activity of glucocorticoids (PubMed:<a href="http://www.uniprot.org/citations/15908412" target="\_blank">15908412</a>, PubMed:<a href="http://www.uniprot.org/citations/17443469" target="\_blank">17443469</a>, PubMed:<a href="http://www.uniprot.org/citations/23776208" target="\_blank">23776208</a>). Has phenylpyruvate tautomerase and dopachrome tautomerase activity (in vitro), but the physiological substrate is not known (PubMed:<a href="http://www.uniprot.org/citations/11439086" target="\_blank">11439086</a>, PubMed:<a href="http://www.uniprot.org/citations/17526494" target="\_blank">17526494</a>). It is not clear whether the tautomerase activity has any physiological relevance, and whether it is important for cytokine activity (PubMed:<a href="http://www.uniprot.org/citations/11439086" target="\_blank">11439086</a>, PubMed:<a href="http://www.uniprot.org/citations/17526494" target="\_blank">17526494</a>).

**Cellular Location**

Secreted. Cytoplasm. Note=Does not have a cleavable signal sequence and is secreted via a specialized, non-classical pathway Secreted by macrophages upon stimulation by bacterial lipopolysaccharide (LPS), or by M.tuberculosis antigens

**Anti-Macrophage migration inhibitory factor (MIF) (CHICKEN) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Anti-Macrophage migration inhibitory factor (MIF) (CHICKEN) Antibody - Images**



Western blot analysis of ROCKLAND Immunochemical's IgY fraction of Chicken anti-Human MIF polyclonal antibody. Lane 2 shows the detection of 100 ng of recombinant MIF present with a single band at 12.3 kDa. Similar detection of MIF will occur when human serum is analyzed. In lane 1 no reaction is observed in the control. A 4-20% gradient gel was used to separate the proteins by SDS-PAGE. The protein was transferred to nitro-cellulose using standard methods. After blocking the membrane was probed with the primary antibody for 1 h at room temperature followed by washes and reaction with a 1:5,000 dilution of IRDye™ 800 conjugated Gt-a-Chicken Rabbit IgG [H&L] (code 603-132-126) for 1 h at room temperature. LICOR's Odyssey® Infrared Imaging System was used to scan and process the image. Other detection systems will yield similar results.

#### **Anti-Macrophage migration inhibitory factor (MIF) (CHICKEN) Antibody - Background**

Macrophage migration inhibitory factor (MIF) is a pro-inflammatory cytokine. It is involved in the innate immune response to bacterial pathogens. The expression of MIF at sites of inflammation suggests a role as mediator in regulating the function of macrophages in host defense. It counteracts the anti-inflammatory activity of glucocorticoids. MIF has phenylpyruvate tautomerase and dopachrome tautomerase activity (in vitro), but the physiological substrate is not known. It is not clear whether the tautomerase activity has any physiological relevance, and whether it is important for cytokine activity.