

**Anti-TSLP (RABBIT) Antibody**  
**TSLP Antibody**  
**Catalog # ASR4941****Specification**

---

**Anti-TSLP (RABBIT) Antibody - Product Information**

Host	<b>Rabbit</b>
Conjugate	<b>Unconjugated</b>
Target Species	<b>Human</b>
Reactivity	<b>Human</b>
Clonality	<b>Polyclonal</b>
Application	<b>WB, E, I, LCI</b>
Application Note	<b>TSLP purified antibody has been tested in western blotting and suitable for ELISA. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 18 kDa in size corresponding to the human TSLP protein by western blotting in appropriate cell lysate or extract.</b>
Physical State	<b>Liquid (sterile filtered)</b>
Buffer	<b>0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>
Immunogen	<b>Anti-TSLP antibody was prepared from rabbit antiserum after repeated immunizations with a truncated form of recombinant human TSLP protein from E. coli.</b>
Preservative	<b>0.01% (w/v) Sodium Azide</b>

**Anti-TSLP (RABBIT) Antibody - Additional Information****Gene ID** 85480**Other Names**  
85480**Purity**

Anti-TSLP is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. This antibody is specific for human TSLP protein. A BLAST analysis was used to suggest cross-reactivity with TSLP from human sources based on 100% homology with the immunizing sequence. Cross-reactivity with TSLP from other sources has not been determined.

**Storage Condition**

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

### Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

## Anti-TSLP (RABBIT) Antibody - Protein Information

**Name** TSLP

### Function

[Isoform 1]: Cytokine that induces the release of T-cell- attracting chemokines from monocytes and, in particular, enhances the maturation of CD11c(+) dendritic cells. Can induce allergic inflammation by directly activating mast cells.

### Cellular Location

Secreted.

### Tissue Location


Isoform 1 is expressed in a number of tissues including heart, liver and prostate. Isoform 2 is the predominant form in keratinocytes of oral mucosa, skin and in salivary glands. It is secreted into saliva.

## Anti-TSLP (RABBIT) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-TSLP (RABBIT) Antibody - Images



Western Blot of Rabbit anti-TSLP antibody. Lane 1: Human TSLP Recombinant Protein. Load: 50 ng per lane. Primary antibody: TSLP antibody at 1:1,000 for overnight at 4°C. Secondary antibody: DyLight™ 649 rabbit secondary antibody (611-143-002) at 1:20,000 for 30 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 13 kDa, 13 kDa for Human TSLP. Other

band(s): None.

### **Anti-TSLP (RABBIT) Antibody - Background**

Thymic Stromal Lymphopoietin (TSLP) is a hematopoietic cytokine. TSLP is mainly expressed in the heart, liver and prostate. TSLP sends signals through a heterodimeric receptor complex, containing the TSLP-R and the IL-7R $\alpha$  chain to induce the release of T cell-attracting chemokines from monocytes. It can induce allergic inflammation by directly activating mast cells. It also promotes T helper type 2 (TH2) cell responses associated with immunity, including atopic dermatitis and asthma, and can promote TH2 cytokine-associated inflammation by directly promoting the effector functions of CD4+ TH2 cells, basophils and other granulocyte populations while simultaneously limiting the expression of dendritic cells derived proinflammatory cytokines and promoting regulatory T cell responses in peripheral tissues. Recombinant human TSLP is a non-glycosylated protein with a molecular weight of 15 kDa. Anti-TSLP antibody is ideal for researchers interested in cytokines and stem cell research.