

Anti-VEGF-165 (RABBIT) Antibody Peroxidase Conjugated
VEGF Antibody Peroxidase Conjugated
Catalog # ASR4960

Specification

Anti-VEGF-165 (RABBIT) Antibody Peroxidase Conjugated - Product Information

Host	Rabbit
Conjugate	Peroxidase (Horseradish)
Target Species	Human
Reactivity	Human
Clonality	Polyclonal
Application	WB, IHC, I, LCI
Application Note	This protein-A purified antibody has been tested for use in western blotting. Reactivity in other assays has not been determined. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 22 kDa in size corresponding to monomeric human VEGF-165 protein by western blotting in the appropriate cell lysate or extract.
Physical State	Lyophilized
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	This purified antibody was prepared from whole rabbit serum produced by repeated immunizations with full length recombinant human VEGF-165 protein.
Reconstitution Volume	100 µL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Anti-VEGF-165 (RABBIT) Antibody Peroxidase Conjugated - Additional Information

Gene ID 7422

Other Names
7422

Purity

This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. This purified antibody has been heated to 56°C for 30 minutes. In ELISA and other immunoreactive assays, this antibody will recognize both native and recombinant human VEGF-165 in cell supernatants and certain body fluids. A control of similarly diluted normal rabbit IgG is recommended.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-VEGF-165 (RABBIT) Antibody Peroxidase Conjugated - Protein Information

Name VEGFA

Synonyms VEGF

Function

[N-VEGF]: Participates in the induction of key genes involved in the response to hypoxia and in the induction of angiogenesis such as HIF1A (PubMed:35455969). Involved in protecting cells from hypoxia-mediated cell death (By similarity).

Cellular Location

[N-VEGF]: Cytoplasm. Nucleus. Note=Cytoplasmic in normoxic conditions and localizes to the nucleus under hypoxic conditions [Isoform L-VEGF189]: Endoplasmic reticulum. Golgi apparatus. Secreted, extracellular space, extracellular matrix [Isoform VEGF165]: Secreted


Tissue Location


Higher expression in pituitary tumors than the pituitary gland. [Isoform VEGF165]: Widely expressed. [Isoform VEGF206]: Not widely expressed.

Anti-VEGF-165 (RABBIT) Antibody Peroxidase Conjugated - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-VEGF-165 (RABBIT) Antibody Peroxidase Conjugated - Images



Western Blot showing detection of Recombinant Human VEGF-165. 50ng of protein (Lane 1) was run on a 4-20% gel and transferred to 0.45 μ m nitrocellulose. After blocking with 1% BSA-TTBS (p/n MB-013, diluted to 1X) 30 min at 20°C, Anti-VEGF-165 (RABBIT) Antibody Peroxidase Conjugate (p/n 209-403-B99) secondary antibody was used at 1:1000 in Blocking Buffer for Fluorescent Western Blotting (p/n MB-070) and imaged using the Bio-Rad VersaDoc® 4000 MP. Arrow indicates correct 19 kDa molecular weight position expected for rH-VEGF-165.

Anti-VEGF-165 (RABBIT) Antibody Peroxidase Conjugated - Background

VEGF (Vascular Endothelial Growth Factor A) is a homodimeric, disulfide-linked glycoprotein involved in angiogenesis which promotes tumor progression and metastasis. It exhibits potent mitogenic and permeability inducing properties specific for the vascular endothelium. Of the four isoforms of VEGF, the smaller two, VEGF 165 and VEGF 121, are secreted proteins and act as diffusible agents, whereas the larger two (VEGF 189 and VEGF 206) remain cell associated. The sequence of this isoform differs from the canonical sequence as follows: 141-141: K \rightarrow N and 142-182: missing. This isoform is often found as a disulfide linked homodimer.