

**Anti-Apolipoprotein C-I (GOAT) Antibody**  
**APOLIPOPROTEIN C-I Antibody**  
Catalog # ASR5071**Specification**

---

**Anti-Apolipoprotein C-I (GOAT) Antibody - Product Information**

Host	Goat
Conjugate	Unconjugated
Target Species	Human
Reactivity	Human
Clonality	Polyclonal
Application	WB, IHC, E, IP, I, LCI
Application Note	Anti-Apolipoprotein antibodies have been used for indirect trapping ELISA for quantitation of antigen in serum using a standard curve, for immunoprecipitation and for western blotting for highly sensitive qualitative analysis.
Physical State	Liquid (sterile filtered)
Buffer	0.125 M Sodium Borate, 0.075 M Sodium Chloride, 0.005 M EDTA, pH 8.0
Immunogen	apoLipoprotein Type C-I produced synthetically in full-length form (not selected epitopes) using conventional peptide technology.
Preservative	0.01% (w/v) Sodium Azide

**Anti-Apolipoprotein C-I (GOAT) Antibody - Additional Information****Gene ID** 341**Other Names**  
341**Purity**

This product has been prepared by immunoaffinity chromatography using immobilized antigens followed by extensive cross-adsorption against other apoLipoproteins and human serum proteins to remove any unwanted specificities. Typically less than 1% cross reactivity against other types of apoLipoprotein was detected by ELISA against purified standards. This antibody reacts with human apoLipoprotein C-I and has negligible cross-reactivity with Type A-I, A-II, B, C-II, C-III, E and J apoLipoproteins. Specific cross reaction of anti-apoLipoprotein antibodies with antigens from other species has not been determined. Non-specific cross reaction of anti-apoLipoprotein antibodies with other human serum proteins is negligible.

**Storage Condition**

Store vial at 4° C prior to opening. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage, mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

## **Anti-Apolipoprotein C-I (GOAT) Antibody - Protein Information**

**Name** APOC1

### **Function**

Inhibitor of lipoprotein binding to the low density lipoprotein (LDL) receptor, LDL receptor-related protein, and very low density lipoprotein (VLDL) receptor. Associates with high density lipoproteins (HDL) and the triacylglycerol-rich lipoproteins in the plasma and makes up about 10% of the protein of the VLDL and 2% of that of HDL. Appears to interfere directly with fatty acid uptake and is also the major plasma inhibitor of cholesteryl ester transfer protein (CETP). Binds free fatty acids and reduces their intracellular esterification. Modulates the interaction of APOE with beta-migrating VLDL and inhibits binding of beta-VLDL to the LDL receptor-related protein.

### **Cellular Location**

Secreted.

### **Tissue Location**

Synthesized mainly in liver and to a minor degree in intestine. Also found in the lung and spleen

## **Anti-Apolipoprotein C-I (GOAT) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Anti-Apolipoprotein C-I (GOAT) Antibody - Images**

## **Anti-Apolipoprotein C-I (GOAT) Antibody - Background**

This antibody is suitable for cardiovascular research.