

**Anti-V5 EPI TOPE TAG (RABBIT) Antibody**  
**V5 Epitope Tag Antibody**  
**Catalog # ASR5193****Specification**

---

**Anti-V5 EPI TOPE TAG (RABBIT) Antibody - Product Information**

Host	Rabbit
Conjugate	Unconjugated
Clonality	Polyclonal
Application	WB, IHC, E, I, LCI
Application Note	Anti-V5 is optimally suited for monitoring expression of V5-tagged fusion proteins. The V5 epitope tag is derived from a small epitope (Pk) present on the P and V proteins of the paramyxovirus of simian virus 5 (SV5). The V5 tag is usually used with all 14 amino acids (GKPIP NPLLGLDST), although it has also been used with a shorter 9 amino acid sequence (IPNPLLGLD). This antibody has been tested by ELISA and western blotting against both the immunizing peptide and V5 containing recombinant proteins. Although not tested, this antibody is likely functional for immunoprecipitation and immunocytochemistry.
Physical State	Liquid (sterile filtered)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	This antibody was purified from whole rabbit serum prepared by repeated immunizations with V5 epitope tag peptide corresponding to aa 95-108 of the V protein conjugated to KLH using maleimide.
Preservative	0.01% (w/v) Sodium Azide

**Anti-V5 EPI TOPE TAG (RABBIT) Antibody - Additional Information****Purity**

This affinity-purified antibody is directed against V5 motif and is useful in determining its presence in various assays. This polyclonal anti-V5-tag antibody detects over-expressed proteins containing the V5 epitope tag. To date this antibody has reacted with all V5 tagged proteins tested so far. In western blotting of bacterial extracts the antibody does not cross-react with endogenous proteins. The antibody recognizes the V5-epitope tag (GKPIP NPLLGLDST) fused to either the carboxy-terminal end of targeted proteins in transfected or transformed cells. Although not yet tested, expect reactivity with recombinant proteins prepared with the V5-epitope tag fused to the amino terminal end as well.

**Storage Condition**

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

#### Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.


### Anti-V5 EPI TOPE TAG (RABBIT) Antibody - Protein Information

### Anti-V5 EPI TOPE TAG (RABBIT) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-V5 EPI TOPE TAG (RABBIT) Antibody - Images



Anti-V5 epitope tag polyclonal antibody detects V5-tagged recombinant protein by western blot. This antibody was used at 1.0 µg/ml to detect 0.05 µg (lane 2) of full-length recombinant mouse serum albumin containing the V5 epitope tag at the carboxy end. Comparison to MW markers (lane 1) indicates detection of monomeric V5 tagged albumin. A 4-20% gradient gel was used to separate the protein by SDS-PAGE under non-reducing conditions. The protein was transferred to nitrocellulose using standard methods. After blocking the membrane was probed with the primary antibody overnight at 4° C followed by washes and reaction with a 1:10,000 dilution of IRDye 800 conjugated Gt-a-Rabbit IgG [H&L] (code 611-132-122) for 45 min at room temperature. LICOR's Odyssey® Infrared Imaging System was used to scan and process the image. Other detection systems will yield similar results.

### Anti-V5 EPI TOPE TAG (RABBIT) Antibody - Background

Epitope tags are short peptide sequences that are easily recognized by tag-specific antibodies.

Due to their small size, epitope tags do not affect the tagged protein's biochemical properties. Most often sequences encoding the epitope tag are included with target DNA at the time of cloning to produce fusion proteins containing the epitope tag sequence. This allows anti-epitope tag antibodies to serve as universal detection reagents for any tag containing protein produced by recombinant means. This means that anti-epitope tag antibodies are a useful alternative to generating specific antibodies to identify, immunoprecipitate or immunoaffinity purify a recombinant protein. The anti-epitope tag antibody is usually functional in a variety of antibody-dependent experimental procedures. Expression vectors producing epitope tag fusion proteins are available for a variety of host expression systems including bacteria, yeast, insect and mammalian cells. Rockland Immunochemicals produces anti-epitope tag antibodies against many common epitope tags including Myc, GST, GFP, 6X His, MBP, FLAG, HA and V5. Rockland Immunochemicals also produces antibodies to other tags including FITC, Rhodamine (TRITC), DNP and biotin.