

**Anti-Protein Kinase C beta (RABBIT) Antibody**  
**Protein Kinase C Beta Antibody**  
**Catalog # ASR5427****Specification****Anti-Protein Kinase C beta (RABBIT) Antibody - Product Information**

|                  |   |
|------------------|---|
| Host             | Rabbit  |
| Conjugate        | Unconjugated  |
| Target Species   | Human   |
| Reactivity       | Human   |
| Clonality        | Polyclonal  |
| Application      | WB, E, I, LCI   |
| Application Note | This affinity purified antibody has been tested for use in ELISA and western blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 76 kDa in size corresponding to PKC beta by western blotting in the appropriate cell lysate or extract. Lysates from U251 or BxPC3 cells are suggested for western blotting. |
| Physical State   | Liquid (sterile filtered)   |
| Buffer           | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |
| Immunogen        | This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region of human Protein kinase C beta.  |
| Preservative     | 0.01% (w/v) Sodium Azide  |

**Anti-Protein Kinase C beta (RABBIT) Antibody - Additional Information****Gene ID** 5579**Other Names**  
5579**Purity**

This product was affinity purified from monospecific antiserum by immunoaffinity chromatography. This antibody is specific for human PKC beta protein. A BLAST analysis was used to suggest cross-reactivity with PKC beta from human and monkey sources based on a 100% homology with the immunizing sequence. Expect partial cross reactivity to PKC beta from rabbit and bovine (93% homology) and rat and mouse (87% homology) sources. Cross-reactivity with PKC beta from other sources has not been determined.

**Storage Condition**

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after

standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

### Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

## Anti-Protein Kinase C beta (RABBIT) Antibody - Protein Information

**Name** PRKCB

**Synonyms** PKCB, PRKCB1

### Function

Calcium-activated, phospholipid- and diacylglycerol (DAG)- dependent serine/threonine-protein kinase involved in various cellular processes such as regulation of the B-cell receptor (BCR) signalosome, oxidative stress-induced apoptosis, androgen receptor-dependent transcription regulation, insulin signaling and endothelial cells proliferation. Plays a key role in B-cell activation by regulating BCR- induced NF-kappa-B activation. Mediates the activation of the canonical NF-kappa-B pathway (NFKB1) by direct phosphorylation of CARD11/CARMA1 at 'Ser-559', 'Ser-644' and 'Ser-652'. Phosphorylation induces CARD11/CARMA1 association with lipid rafts and recruitment of the BCL10-MALT1 complex as well as MAP3K7/TAK1, which then activates IKK complex, resulting in nuclear translocation and activation of NFKB1. Plays a direct role in the negative feedback regulation of the BCR signaling, by down-modulating BTK function via direct phosphorylation of BTK at 'Ser-180', which results in the alteration of BTK plasma membrane localization and in turn inhibition of BTK activity (PubMed:<a href="http://www.uniprot.org/citations/11598012" target="\_blank">11598012</a>). Involved in apoptosis following oxidative damage: in case of oxidative conditions, specifically phosphorylates 'Ser-36' of isoform p66Shc of SHC1, leading to mitochondrial accumulation of p66Shc, where p66Shc acts as a reactive oxygen species producer. Acts as a coactivator of androgen receptor (AR)-dependent transcription, by being recruited to AR target genes and specifically mediating phosphorylation of 'Thr-6' of histone H3 (H3T6ph), a specific tag for epigenetic transcriptional activation that prevents demethylation of histone H3 'Lys-4' (H3K4me) by LSD1/KDM1A (PubMed:<a href="http://www.uniprot.org/citations/20228790" target="\_blank">20228790</a>). In insulin signaling, may function downstream of IRS1 in muscle cells and mediate insulin-dependent DNA synthesis through the RAF1-MAPK/ERK signaling cascade. Participates in the regulation of glucose transport in adipocytes by negatively modulating the insulin-stimulated translocation of the glucose transporter SLC2A4/GLUT4. Phosphorylates SLC2A1/GLUT1, promoting glucose uptake by SLC2A1/GLUT1 (PubMed:<a href="http://www.uniprot.org/citations/25982116" target="\_blank">25982116</a>). Under high glucose in pancreatic beta-cells, is probably involved in the inhibition of the insulin gene transcription, via regulation of MYC expression. In endothelial cells, activation of PRKCB induces increased phosphorylation of RB1, increased VEGFA-induced cell proliferation, and inhibits PI3K/AKT-dependent nitric oxide synthase (NOS3/eNOS) regulation by insulin, which causes endothelial dysfunction. Also involved in triglyceride homeostasis (By similarity). Phosphorylates ATF2 which promotes cooperation between ATF2 and JUN, activating transcription (PubMed:<a href="http://www.uniprot.org/citations/19176525" target="\_blank">19176525</a>). Phosphorylates KLHL3 in response to angiotensin II signaling, decreasing the interaction between KLHL3 and WNK4 (PubMed:<a href="http://www.uniprot.org/citations/25313067" target="\_blank">25313067</a>). Phosphorylates and activates LRRK1, which phosphorylates RAB proteins involved in intracellular trafficking (PubMed:<a href="http://www.uniprot.org/citations/36040231" target="\_blank">36040231</a>).

### Cellular Location


Cytoplasm. Nucleus. Membrane; Peripheral membrane protein

## Anti-Protein Kinase C beta (RABBIT) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Protein Kinase C beta (RABBIT) Antibody - Images



Western blot using Rockland's affinity purified anti-PKC beta antibody shows detection of PKC beta in ~25 µg of U251 whole cell lysate (glioma derived). A 4-12% gel was used for separation. The arrowhead corresponds to 76 kDa PKC beta. The membrane was probed with the primary antibody at a 1:1,000 dilution in 5% BSA in TTBS at 4° C, overnight. Personal Communication, Howard Fine and Svetlana Kotliarova, CCR-NCI, Bethesda, MD.

## Anti-Protein Kinase C beta (RABBIT) Antibody - Background

This antibody is designed, produced, and validated as part of a collaboration between Rockland and the National Cancer Institute (NCI) and is suitable for Cancer, Immunology and Nuclear Signaling research. Protein kinase C (PKC) beta is implicated as a potential important factor in cancer cell proliferation and angiogenesis. PKC is a family of serine- and threonine-specific protein kinases that can be activated by calcium and second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This protein kinase has been reported to be involved in many different cellular functions, such as B cell activation, apoptosis induction, endothelial cell proliferation, and intestinal sugar absorption. Studies in mice also suggest that this kinase may also regulate neuronal functions and correlate fear-induced conflict behavior after stress. Alternatively spliced transcript variants encoding distinct isoforms have been reported.