

**Anti-GAB1 (RABBIT) Antibody**  
**GAB1 Antibody**  
**Catalog # ASR5745**

**Specification**

---

**Anti-GAB1 (RABBIT) Antibody - Product Information**

Host	Rabbit
Conjugate	Unconjugated
Target Species	Human
Reactivity	Human
Clonality	Polyclonal
Application	WB, E, I, LCI
Application Note	Anti-GAB1 Antibody has been tested for use in ELISA and by western blot. Specific conditions for reactivity should be optimized by the end user. Expect a predominant band at 76.6 kDa by western blotting in the appropriate cell lysate or extract.
Physical State	Liquid (sterile filtered)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Anti-GAB1 Antibody was produced in rabbits by repeated immunizations with a synthetic peptide corresponding to residues surrounding Y659 of human GAB1 protein.
Preservative	0.01% (w/v) Sodium Azide

**Anti-GAB1 (RABBIT) Antibody - Additional Information**

**Gene ID** 2549

**Other Names**  
2549

**Purity**

Anti-GAB1 was prepared from monospecific antiserum by immunoaffinity chromatography using phospho peptide coupled to agarose beads followed by solid phase adsorptions against non-phospho peptide and non-specific peptide to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum. This antibody is specific for human GAB1. Cross-reactivity against GAB1 from other species may occur but has not yet been tested.

**Storage Condition**

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**Anti-GAB1 (RABBIT) Antibody - Protein Information**

**Name** GAB1


**Function**

Adapter protein that plays a role in intracellular signaling cascades triggered by activated receptor-type kinases. Plays a role in FGFR1 signaling. Probably involved in signaling by the epidermal growth factor receptor (EGFR) and the insulin receptor (INSR). Involved in the MET/HGF-signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/29408807" target="\_blank">29408807</a>).

**Anti-GAB1 (RABBIT) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-GAB1 (RABBIT) Antibody - Images**

Western Blot of Rabbit anti-GAB1 antibody. Left Blot: Phospho GAB1 rProtein. Right Blot: Phospho GAB1 rProtein incubated with GAB1 immunizing peptide. Load: 0.05  $\mu$ g per lane. Primary antibody: GAB1 antibody at 1  $\mu$ g/mL for overnight at 4°C. Secondary antibody: HRP Goat anti-Rabbit IgG secondary antibody (p/n 611-103-122) at 1:40,000 for 45 min at RT. Block: (p/n MB-070) Fluorescent blocking buffer overnight at 4°C. Predicted/Observed size: 130 kDa for GAB1. Other band(s): none.

**Anti-GAB1 (RABBIT) Antibody - Background**

GAB1 antibody detects human GAB1. GAB1 is a member of the IRS1-like multisubstrate docking

protein family. The protein is an important mediator of branching tubulogenesis and plays a central role in cellular growth response, transformation and apoptosis. Two transcript variants encoding different isoforms have been found for this gene. GAB1 plays a role in intracellular signaling cascades triggered by activated receptor-type kinases. It is known to play a role in FGFR1 signaling and is probably involved in signaling by the epidermal growth factor receptor (EGFR) and the insulin receptor (INSR). GAB1 interacts with GRB2 and with other SH2-containing proteins. It is known to interact with phosphorylated LAT2, PTPRJ, FRS2, GRB2, PIK3R1 and SOS1. GAB1 gets phosphorylated in response to FGFR1 activation. This tyrosine phosphorylation of GAB1 mediates interaction with several proteins that contain SH2 domains. Anti-GAB1 Antibody is ideal for investigators involved in Cell Signaling, Cancer, Neuroscience and Signal Transduction research.