

Anti-Collagen Type III (RABBIT) Antibody Peroxidase Conjugated

Collagen Type III Antibody Peroxidase Conjugated Catalog # ASR5805

Specification

Anti-Collagen Type III (RABBIT) Antibody Peroxidase Conjugated - Product Information

Host Rabbit

Conjugate Peroxidase (Horseradish)

Target Species Mammalian
Reactivity Human, Bovine
Clonality Polyclonal

Application WB, IHC, E, IP, I, LCI

Application Note

Anti-Collagen Type III Peroxidase
Conjugated Antibody is suitable for

western blotting, IHC and for ELISA.
Researchers should determine optimal
titers for applications that are not stated

below. Lvophilized

Physical State Lyophilized

Buffer 0.01 M Sodium Phosphate, 0.25 M Sodium

Chloride, pH 7.2

Immunogen Collagen Type III from human and bovine

placenta

Reconstitution Volume 50 μL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Thimerosal

Anti-Collagen Type III (RABBIT) Antibody Peroxidase Conjugated - Additional Information

Gene ID 1281

Other Names 1281

Purity

This product has been prepared by immunoaffinity chromatography using immobilized antigens. Some class-specific anti-collagens may be specific for three-dimensional epitopes which may result in diminished reactivity with denatured collagen or formalin-fixed, paraffin embedded tissues. This antibody reacts with most mammalian Type III collagens and has expected cross-reactivity with Type I and negligible cross reactivity with Type II, IV, V or VI collagens. Non-specific cross-reaction of anti-collagen antibodies with other human serum proteins or non-collagen extracellular matrix proteins has not been tested.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted



liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Collagen Type III (RABBIT) Antibody Peroxidase Conjugated - Protein Information

Name COL3A1

Function

Collagen type III occurs in most soft connective tissues along with type I collagen. Involved in regulation of cortical development. Is the major ligand of ADGRG1 in the developing brain and binding to ADGRG1 inhibits neuronal migration and activates the RhoA pathway by coupling ADGRG1 to GNA13 and possibly GNA12.

Cellular Location

Secreted, extracellular space, extracellular matrix {ECO:0000255|PROSITE-ProRule:PRU00793}

Anti-Collagen Type III (RABBIT) Antibody Peroxidase Conjugated - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Collagen Type III (RABBIT) Antibody Peroxidase Conjugated - Images

Anti-Collagen Type III (RABBIT) Antibody Peroxidase Conjugated - Background

In muscle tissue, collagen serves as a major component of the endomysium. Collagen constitutes one to two percent of muscle tissue, and accounts for 6% of the weight of strong, tendinous muscles. A collagen may be defined as a protein containing sizable domain(s) of triple-helical conformation. Type IV collagen is a major macromolecular constituent of basement membranes and can be readily isolated from basement-membrane-rich tissues or highly vascularized tissues such as the placental villi. This collagen appears to be largely restricted to structures identifiable as basement membranes. In contrast, type VI collagen appears to be prevalent in several tissues even though it has been isolated largely from placental villi preparations. The extent to which type VII and VIII collagens are distributed is not known.