

**Anti-Collagen Type VI (RABBIT) Antibody Peroxidase Conjugated**  
**Collagen Type VI Antibody Peroxidase Conjugated**  
**Catalog # ASR5808****Specification****Anti-Collagen Type VI (RABBIT) Antibody Peroxidase Conjugated - Product Information**

Host	Rabbit
Conjugate	Peroxidase (Horseradish)
Target Species	Mammalian
Reactivity	Human, Bovine
Clonality	Polyclonal
Application	WB, IHC, E, IP, I, LCI
Application Note	Anti-Collagen Type VI (RABBIT) Peroxidase Conjugated Antibody is suitable for western blot, immunoprecipitation, and immunohistochemistry. Researchers should determine optimal titers for applications that are not stated below.
Physical State	Lyophilized
Buffer	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Immunogen	Collagen Type VI from human and bovine placenta
Reconstitution Volume	50 µL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Thimerosal

**Anti-Collagen Type VI (RABBIT) Antibody Peroxidase Conjugated - Additional Information****Gene ID** 1291**Other Names**

1291

**Purity**

Anti-Collagen VI has been prepared by immunoaffinity chromatography using immobilized antigens followed by extensive cross-adsorption against other collagens, human serum proteins and non-collagen extracellular matrix proteins to remove any unwanted specificities. Some class specific anti-collagens may be specific for three-dimensional epitopes which may result in diminished reactivity with denatured collagen or formalin-fixed, paraffin embedded tissues. This antibody reacts with most mammalian Type VI collagens and has negligible cross-reactivity with Type I, II, III, IV and V collagens. Non-specific cross reaction of anti-collagen antibodies with other human serum proteins or non-collagen extracellular matrix proteins is negligible.

**Storage Condition**

Store vial at 4° C prior to restoration. Restore with 0.05 mL of deionized water (or equivalent). For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and

thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**Anti-Collagen Type VI (RABBIT) Antibody Peroxidase Conjugated - Protein Information**

**Name** COL6A1

**Function**

Collagen VI acts as a cell-binding protein.

**Cellular Location**

Secreted, extracellular space, extracellular matrix

**Anti-Collagen Type VI (RABBIT) Antibody Peroxidase Conjugated - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-Collagen Type VI (RABBIT) Antibody Peroxidase Conjugated - Images****Anti-Collagen Type VI (RABBIT) Antibody Peroxidase Conjugated - Background**

Collagen Type VI Peroxidase Conjugated Antibody is specific for collagen type VI protein. The collagens are a superfamily of proteins that play a role in maintaining the integrity of various tissues. Collagen VI is a major structural component of microfibrils. The basic structural unit of collagen VI is a heterotrimer of the alpha1(VI), alpha2(VI), and alpha3(VI) chains. The alpha2(VI) and alpha3(VI) chains are encoded by the COL6A2 and COL6A3 genes, respectively. The protein encoded by this gene is the alpha 1 subunit of type VI collagen (alpha1(VI) chain). Mutations in the genes that code for the collagen VI subunits result in the autosomal dominant disorder, Bethlem myopathy.