

**Anti-ATM Protein Kinase pS1981 (SHEEP) Antibody**  
**ATM phospho S1981 Antibody**  
**Catalog # ASR5896****Specification**

---

**Anti-ATM Protein Kinase pS1981 (SHEEP) Antibody - Product Information**

Host	Sheep
Conjugate	Unconjugated
Target Species	Human
Reactivity	Human
Clonality	Polyclonal
Application	WB, E, I, LCI
Application Note	Affinity purified sheep anti-ATM has been tested by ELISA and western blotting against native form phosphorylated ATM PK.
Physical State	Liquid (sterile filtered)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	This antibody was affinity purified from whole sheep serum prepared by repeated immunizations with a synthetic peptide corresponding to a region near serine 1981 of human ATM conjugated to KLH using maleimide.
Preservative	0.01% (w/v) Sodium Azide

**Anti-ATM Protein Kinase pS1981 (SHEEP) Antibody - Additional Information****Gene ID** 472**Other Names**  
472**Purity**

This affinity-purified antibody is directed against human ATM and is useful in determining its presence in various assays. This polyclonal anti-ATM antibody recognizes the phosphorylated epitope in native and over-expressed proteins found in various tissues and extracts. Reactivity is observed against human ATM and cross reactivity with ATM from other mammalian sources has not been tested.

**Storage Condition**

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.



[21757780](http://www.uniprot.org/citations/21757780), PubMed:<[24534091](http://www.uniprot.org/citations/24534091)>, PubMed:<[26240375](http://www.uniprot.org/citations/26240375)>, PubMed:<[26774286](http://www.uniprot.org/citations/26774286)>, PubMed:<[30612738](http://www.uniprot.org/citations/30612738)>, PubMed:<[30886146](http://www.uniprot.org/citations/30886146)>, PubMed:<[30952868](http://www.uniprot.org/citations/30952868)>, PubMed:<[38128537](http://www.uniprot.org/citations/38128537)>, PubMed:<[9733515](http://www.uniprot.org/citations/9733515)>, PubMed:<[9843217](http://www.uniprot.org/citations/9843217)>). May play a role in vesicle and/or protein transport. Could play a role in T-cell development, gonad and neurological function. Plays a role in replication-dependent histone mRNA degradation. Binds DNA ends. Phosphorylation of DYRK2 in nucleus in response to genotoxic stress prevents its MDM2-mediated ubiquitination and subsequent proteasome degradation (PubMed:<[19965871](http://www.uniprot.org/citations/19965871)>). Phosphorylates ATF2 which stimulates its function in DNA damage response (PubMed:<[15916964](http://www.uniprot.org/citations/15916964)>). Phosphorylates ERCC6 which is essential for its chromatin remodeling activity at DNA double-strand breaks (PubMed:<[29203878](http://www.uniprot.org/citations/29203878)>). Phosphorylates TTC5/STRAP at 'Ser-203' in the cytoplasm in response to DNA damage, which promotes TTC5/STRAP nuclear localization (PubMed:<[15448695](http://www.uniprot.org/citations/15448695)>). Also involved in pexophagy by mediating phosphorylation of PEX5: translocated to peroxisomes in response to reactive oxygen species (ROS), and catalyzes phosphorylation of PEX5, promoting PEX5 ubiquitination and induction of pexophagy (PubMed:<[26344566](http://www.uniprot.org/citations/26344566)>).

#### Cellular Location

Nucleus. Cytoplasmic vesicle. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250|UniProtKB:Q62388}. Peroxisome matrix. Note=Primarily nuclear (PubMed:9050866, PubMed:9150358). Found also in endocytic vesicles in association with beta-adaptin (PubMed:9707615). Translocated to peroxisomes in response to reactive oxygen species (ROS) by PEX5 (PubMed:26344566)

#### Tissue Location


Found in pancreas, kidney, skeletal muscle, liver, lung, placenta, brain, heart, spleen, thymus, testis, ovary, small intestine, colon and leukocytes


### Anti-ATM Protein Kinase pS1981 (SHEEP) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-ATM Protein Kinase pS1981 (SHEEP) Antibody - Images





Western Blot of Sheep Anti-ATM pS1981 polyclonal antibody. Lane 1: untreated MCF-7 cell lysate (p/n W09-000-360). Lane 2: Hydrogen Peroxide stimulated MCF-7 Whole Cell Lysate (p/n W09-000-366). Load: 35  $\mu$ g per lane. Primary antibody: ATM pS1981 antibody at 1:1000 for 1 h at room temperature. Secondary antibody: IRDye™ 800 conjugated Donkey anti-Sheep IgG secondary antibody at 1:5,000 for 1h at room temperature. Block: 5% BLOTTO (B501) overnight at 4°C. Predicted/Observed size: 370 kDa, ATM (370 kDa) is indicated by an arrow. Other band(s): ATM splice variants and isoforms.

#### **Anti-ATM Protein Kinase pS1981 (SHEEP) Antibody - Background**

ATM, the gene mutated in the hereditary disease ataxia-telangiectasia, codes for a protein kinase that acts as a master regulator of cellular responses to DNA double-strand breaks. ATM is normally inactive and the question of how it is activated in the event of DNA damage (due to ionizing radiation for instance) is central to understanding its function. ATM protein is now shown to be present in undamaged cells as an inactive dimer. Low doses of ionizing radiation, which induce only a few DNA breaks, activate at least half of the total ATM protein present, possibly in response to changes in chromatin structure. The ATM gene encodes a 370-kDa protein that belongs to the phosphoinositide 3-kinase (PI(3)K) superfamily, but which phosphorylates proteins rather than lipids. The 350-amino-acid kinase domain at the carboxy terminus of this large protein is the only segment of ATM with an assigned function. Exposure of cells to IR triggers ATM kinase activity and this function is required for arrests in G1, S and G2 phases of the cell cycle. Several substrates of the ATM kinase participate in these IR-induced cell-cycle arrests. These include p53, Mdm2 and Chk2 in the G1 checkpoint; Nbs1, Brca1, FancD2 and SMC1 in the transient IR-induced S-phase arrest; and Brca1 and hRad17 in the G2/M checkpoint. This antibody is similar to the rabbit host antibody discussed by Bakkenist, C. J. & Kastan, M. B. in *Nature* 421, 499-506 (2003).