

ABCC2 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant ABCC2.

Catalog # AT1005a

Specification

ABCC2 Antibody (monoclonal) (M01) - Product Information

Application	WB, E
Primary Accession	O92887
Other Accession	NM_000392
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	174207

ABCC2 Antibody (monoclonal) (M01) - Additional Information

Gene ID 1244

Other Names

Canalicular multispecific organic anion transporter 1, ATP-binding cassette sub-family C member 2, Canalicular multidrug resistance protein, Multidrug resistance-associated protein 2, ABCC2, CMOAT, CMOAT1, CMRP, MRP2

Target/Specificity

ABCC2 (NP_000383, 214 a.a. ~ 313 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

ABCC2 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


ABCC2 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ABCC2 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .



Detection limit for recombinant GST tagged ABCC2 is approximately 0.03ng/ml as a capture antibody.

ABCC2 Antibody (monoclonal) (M01) - Background

The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MRP subfamily which is involved in multi-drug resistance. This protein is expressed in the canalicular (apical) part of the hepatocyte and functions in biliary transport. Substrates include anticancer drugs such as vinblastine; therefore, this protein appears to contribute to drug resistance in mammalian cells. Several different mutations in this gene have been observed in patients with Dubin-Johnson syndrome (DJS), an autosomal recessive disorder characterized by conjugated hyperbilirubinemia.

ABCC2 Antibody (monoclonal) (M01) - References

Influence of CYP3A5 and drug transporter polymorphisms on imatinib trough concentration and clinical response among patients with chronic phase chronic myeloid leukemia. Takahashi N, et al. J Hum Genet, 2010 Aug 19. PMID 20720558. Pharmacogenetic analysis of lipid responses to

rosuvastatin in Chinese patients. Hu M, et al. Pharmacogenet Genomics, 2010 Oct. PMID 20679960. Risk of diarrhoea in a long-term cohort of renal transplant patients given mycophenolate mofetil: the significant role of the UGT1A8 2 variant allele. Woillard JB, et al. Br J Clin Pharmacol, 2010 Jun. PMID 20565459. Pharmacogenetic predictors of adverse events and response to chemotherapy in metastatic colorectal cancer: results from North American Gastrointestinal Intergroup Trial N9741. McLeod HL, et al. J Clin Oncol, 2010 Jul 10. PMID 20530282. A Large-scale genetic association study of esophageal adenocarcinoma risk. Liu CY, et al. Carcinogenesis, 2010 Jul. PMID 20453000.