

**ABHD5 Antibody (monoclonal) (M02)**

Mouse monoclonal antibody raised against a partial recombinant ABHD5.

Catalog # AT1013a

**Specification**

---

**ABHD5 Antibody (monoclonal) (M02) - Product Information**

Application	WB, E
Primary Accession	<a href="#">Q8WTS1</a>
Other Accession	<a href="#">NM_016006</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	39096

**ABHD5 Antibody (monoclonal) (M02) - Additional Information**

Gene ID 51099

**Other Names**

1-acylglycerol-3-phosphate O-acyltransferase ABHD5, Abhydrolase domain-containing protein 5, Lipid droplet-binding protein CGI-58, ABHD5, NCIE2

**Target/Specificity**

ABHD5 (NP\_057090, 240 a.a. ~ 342 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

ABHD5 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.


**ABHD5 Antibody (monoclonal) (M02) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**ABHD5 Antibody (monoclonal) (M02) - Images**



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.07 KDa) .



ABHD5 monoclonal antibody (M02), clone 4B12 Western Blot analysis of ABHD5 expression in A-431 ( (Cat # AT1013a )



Detection limit for recombinant GST tagged ABHD5 is 0.3 ng/ml as a capture antibody.

**ABHD5 Antibody (monoclonal) (M02) - Background**

The protein encoded by this gene belongs to a large family of proteins defined by an alpha/beta hydrolase fold, and contains three sequence motifs that correspond to a catalytic triad found in the esterase/lipase/thioesterase subfamily. It differs from other members of this subfamily in that its putative catalytic triad contains an asparagine instead of the serine residue. Mutations in this gene have been associated with Chanarin-Dorfman syndrome, a triglyceride storage disease with impaired long-chain fatty acid oxidation.

### **ABHD5 Antibody (monoclonal) (M02) - References**

1.The Cannabinoid Receptor Agonist THC Attenuates Weight Loss in a Rodent Model of Activity-Based Anorexia.Verty AN, Evetts MJ, Crouch GJ, McGregor IS, Stefanidis A, Oldfield BJ.Neuropsychopharmacology. 2011 Jun;36(7):1349-58. Epub 2011 Mar 16.2.Adipose triacylglycerol lipase deletion alters whole body energy metabolism and impairs exercise performance in mice.Huijsman E, van de Par C, Economou C, van der Poel C, Lynch GS, Schoiswohl G, Haemmerle G, Zechner R, Watt MJ.Am J Physiol Endocrinol Metab. 2009 Aug;297(2):E505-13. Epub 2009 Jun 2.3.Adipose Triglyceride Lipase Regulation of Skeletal Muscle Lipid Metabolism and Insulin Responsiveness.Watt MJ, van Denderen BJ, Castelli LA, Bruce CR, Hoy AJ, Kraegen EW, Macaulay L, Kemp BE.Mol Endocrinol. 2008 May;22(5):1200-12. Epub 2008 Jan 17.