

ARRB2 Antibody (monoclonal) (M06)

Mouse monoclonal antibody raised against a partial recombinant ARRB2.

Catalog # AT1202a

Specification

ARRB2 Antibody (monoclonal) (M06) - Product Information

Application	IF, WB, E
Primary Accession	P32121
Other Accession	BC007427
Reactivity	Human, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	46106

ARRB2 Antibody (monoclonal) (M06) - Additional Information

Gene ID 409

Other Names

Beta-arrestin-2, Arrestin beta-2, ARRB2, ARB2, ARR2

Target/Specificity

ARRB2 (AAH07427, 300 a.a. ~ 409 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB ~ ~ 1:500 ~ 1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

ARRB2 Antibody (monoclonal) (M06) is for research use only and not for use in diagnostic or therapeutic procedures.


ARRB2 Antibody (monoclonal) (M06) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


ARRB2 Antibody (monoclonal) (M06) - Images




Immunofluorescence of monoclonal antibody to ARRB2 on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.84 kDa) .



ARRB2 monoclonal antibody (M06), clone 4D2. Western Blot analysis of ARRB2 expression in PC-12.



ARRB2 monoclonal antibody (M06), clone 4D2. Western Blot analysis of ARRB2 expression in MCF-7.



Detection limit for recombinant GST tagged ARRB2 is approximately 0.3ng/ml as a capture antibody.

ARRB2 Antibody (monoclonal) (M06) - Background

Members of arrestin/beta-arrestin protein family are thought to participate in agonist-mediated desensitization of G-protein-coupled receptors and cause specific dampening of cellular responses to stimuli such as hormones, neurotransmitters, or sensory signals. Arrestin beta 2, like arrestin beta 1, was shown to inhibit beta-adrenergic receptor function in vitro. It is expressed at high levels in the central nervous system and may play a role in the regulation of synaptic receptors. Besides the brain, a cDNA for arrestin beta 2 was isolated from thyroid gland, and thus it may also be involved in hormone-specific desensitization of TSH receptors. Multiple alternatively spliced transcript variants have been found for this gene, but the full-length nature of some variants has not been defined.

ARRB2 Antibody (monoclonal) (M06) - References

beta-Arrestin2 influences the response to methadone in opioid-dependent patients. Oneda B, et al. *Pharmacogenomics J*, 2010 Jun 1. PMID 20514076. Beta-arrestin 2 regulates Toll-like receptor 4-mediated apoptotic signalling through glycogen synthase kinase-3beta. Li H, et al. *Immunology*, 2010 Aug. PMID 20497256. Preliminary evidence of ethnic divergence in associations of putative genetic variants for methamphetamine dependence. Bousman CA, et al. *Psychiatry Res*, 2010 Jul 30. PMID 20478633. Beta-arrestin 2 is required for B1 receptor-dependent post-translational activation of inducible nitric oxide synthase. Kuhr FK, et al. *FASEB J*, 2010 Jul. PMID 20228252. Non-visual arrestins are constitutively associated with the centrosome and regulate centrosome function. Shankar H, et al. *J Biol Chem*, 2010 Mar 12. PMID 20056609.