

## COX6C Antibody (monoclonal) (M03)

Mouse monoclonal antibody raised against a full length recombinant COX6C.

Catalog # AT1603a

### Specification

---

#### COX6C Antibody (monoclonal) (M03) - Product Information

Application	WB, IHC, E
Primary Accession	<a href="#">P09669</a>
Other Accession	<a href="#">BC000187</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	8781

#### COX6C Antibody (monoclonal) (M03) - Additional Information

Gene ID 1345

#### Other Names

Cytochrome c oxidase subunit 6C, Cytochrome c oxidase polypeptide VIc, COX6C

#### Target/Specificity

COX6C (AAH00187, 1 a.a. ~ 75 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

#### Dilution

WB ~ 1:500 ~ 1000

#### Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

#### Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

#### Precautions

COX6C Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.


#### COX6C Antibody (monoclonal) (M03) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### COX6C Antibody (monoclonal) (M03) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (33.99 KDa) .



Immunoperoxidase of monoclonal antibody to COX6C on formalin-fixed paraffin-embedded human small intestine. [antibody concentration 1 μg/ml]



Detection limit for recombinant GST tagged COX6C is approximately 1 ng/ml as a capture antibody.

### COX6C Antibody (monoclonal) (M03) - Background

Cytochrome c oxidase, the terminal enzyme of the mitochondrial respiratory chain, catalyzes the

electron transfer from reduced cytochrome c to oxygen. It is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may be involved in the regulation and assembly of the complex. This nuclear gene encodes subunit VIc, which has 77% amino acid sequence identity with mouse subunit VIc. This gene is up-regulated in prostate cancer cells. A pseudogene has been found on chromosomes 16p12.

### **COX6C Antibody (monoclonal) (M03) - References**

Assembly of nuclear DNA-encoded subunits into mitochondrial complex IV, and their preferential integration into supercomplex forms in patient mitochondria. Lazarou M, et al. FEBS J, 2009 Nov. PMID 19843159. Polymorphisms in mitochondrial genes and prostate cancer risk. Wang L, et al. Cancer Epidemiol Biomarkers Prev, 2008 Dec. PMID 19064571. Mid-region parathyroid hormone-related protein (PTHrP) and gene expression of MDA-MB231 breast cancer cells. Sirchia R, et al. Biol Chem, 2007 May. PMID 17516841. Large-scale mapping of human protein-protein interactions by mass spectrometry. Ewing RM, et al. Mol Syst Biol, 2007. PMID 17353931. A human protein-protein interaction network: a resource for annotating the proteome. Stelzl U, et al. Cell, 2005 Sep 23. PMID 16169070.