

DDIT3 Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant DDIT3.****Catalog # AT1731a****Specification**

DDIT3 Antibody (monoclonal) (M01) - Product Information

Application	WB, E
Primary Accession	P35638
Other Accession	BC003637
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	19175

DDIT3 Antibody (monoclonal) (M01) - Additional Information**Gene ID** 1649**Other Names**

DNA damage-inducible transcript 3 protein, DDIT-3, C/EBP zeta, C/EBP-homologous protein, CHOP, C/EBP-homologous protein 10, CHOP-10, CCAAT/enhancer-binding protein homologous protein, Growth arrest and DNA damage-inducible protein GADD153, DDIT3, CHOP, CHOP10, GADD153

Target/Specificity

DDIT3 (AAH03637, 1 a.a. ~ 90 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

DDIT3 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


DDIT3 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DDIT3 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (35.53 KDa) .



Detection limit for recombinant GST tagged DDIT3 is approximately 0.03ng/ml as a capture antibody.

DDIT3 Antibody (monoclonal) (M01) - Background

This gene encodes a member of the CCAAT/enhancer-binding protein (C/EBP) family of transcription factors. The protein functions as a dominant-negative inhibitor by forming heterodimers with other C/EBP members, such as C/EBP and LAP (liver activator protein), and preventing their DNA binding activity. The protein is implicated in adipogenesis and erythropoiesis, is activated by endoplasmic reticulum stress, and promotes apoptosis. Fusion of this gene and FUS on chromosome 16 or EWSR1 on chromosome 22 induced by translocation generates chimeric proteins in myxoid liposarcomas or Ewing sarcoma. Multiple alternatively spliced transcript variants encoding two isoforms with different length have been identified.

DDIT3 Antibody (monoclonal) (M01) - References

1. Loss of UDP-N-acetylglucosamine 2-epimerase/ N-acetylmannosamine kinase (GNE) induces apoptotic processes in pancreatic carcinoma cells. Kemmner W, Kessel P, Sanchez-Ruderisch H, Moller H, Hinderlich S, Schlag PM, Detjen K. FASEB J. 2011 Nov 2.