

EXOC3 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant EXOC3.

Catalog # AT1963a

Specification

EXOC3 Antibody (monoclonal) (M01) - Product Information

Application	WB, E
Primary Accession	O60645
Other Accession	NM_007277
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2b Kappa
Calculated MW	85567

EXOC3 Antibody (monoclonal) (M01) - Additional Information

Gene ID 11336

Other Names

Exocyst complex component 3, Exocyst complex component Sec6, EXOC3, SEC6, SEC6L1

Target/Specificity

EXOC3 (NP_009208, 646 a.a. ~ 745 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

EXOC3 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


EXOC3 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EXOC3 Antibody (monoclonal) (M01) - Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .



Western Blot analysis of EXOC3 expression in transfected 293T cell line by EXOC3 monoclonal antibody (M01), clone 4A7.

Lane 1: EXOC3 transfected lysate(86.845 KDa).
 Lane 2: Non-transfected lysate.



Detection limit for recombinant GST tagged EXOC3 is approximately 3ng/ml as a capture antibody.

EXOC3 Antibody (monoclonal) (M01) - Background

The protein encoded by this gene is a component of the exocyst complex, a multiple protein complex essential for targeting exocytic vesicles to specific docking sites on the plasma membrane. Though best characterized in yeast, the component proteins and functions of exocyst complex have been demonstrated to be highly conserved in higher eukaryotes. At least eight components of the exocyst complex, including this protein, are found to interact with the actin cytoskeletal remodeling and vesicle transport machinery. The complex is also essential for the biogenesis of epithelial cell surface polarity.

EXOC3 Antibody (monoclonal) (M01) - References

1. West Nile virus and Dengue virus capsid protein negates the antiviral activity of human Sec3 protein Through The Proteasome Pathway. Raghavan B, Ng M. Cell Microbiol. 2013 Mar 22. doi: 10.1111/cmi.12143. 2. Exocyst subunits are involved in isoproterenol-induced amylase release from rat parotid acinar cells. Imai A, Yoshie S, Haga-Tsujimura M, Nashida T, Shimomura H. Eur J Oral Sci. 2012 Apr;120(2):123-31. doi: 10.1111/j.1600-0722.2012.00952.x.